Wright State University CORE Scholar

Unintentional Prescription Drug Poisoning Project

Center for Interventions, Treatment and Addictions
Research

2011

2011 Poisoning Death Review Narrative Summary

Center for Interventions, Treatment & Addictions Research

Follow this and additional works at: http://corescholar.libraries.wright.edu/prescription_drugs

Part of the <u>Social and Behavioral Sciences Commons</u>

Repository Citation

(2011). 2011 Poisoning Death Review Narrative Summary. . http://corescholar.libraries.wright.edu/prescription_drugs/2

This Report is brought to you for free and open access by the Center for Interventions, Treatment and Addictions Research at CORE Scholar. It has been accepted for inclusion in Unintentional Prescription Drug Poisoning Project by an authorized administrator of CORE Scholar. For more information, please contact corescholar@www.libraries.wright.edu.

2011 Poisoning Death Review Narrative Summary

There were 130 unintentional drug overdose deaths in Montgomery County in 2011. The designation of unintentional (accident), as compared to intentional (suicide), is made by the Montgomery County Coroner's Office. The Poison Death Review (PDR) is the compilation, presentation, and interpretation of data from multiple data streams within the Montgomery County Coroner's Office. This is the second year a PDR has been completed for Montgomery County. The first time was in 2010.

Of the 130 deaths in 2011, 116 were residents of Montgomery County, 6 were residents of other Ohio counties, 6 were out-of-state residents, and 2 individuals had no known address. As in 2010, the age groups most represented were 35-44 year olds and 44-55 year olds, accounting for 58% of the decedents compared to 46% in 2010. Persons aged 25-34 years old accounted for 22% of the decedents. The majority of the decedents were male (59%) and white (87%). In terms of education, about 97% had a high school education or less.

Most decedents, 74%, had a history of physical illness or disability, and 56% had a history of some form of heart disease. A large majority (82%) had a history of substance abuse. Most of the deaths occurred in the decedent's home (53%).

Opioids (prescription or illicit) were the drugs most frequently mentioned in toxicology reports. They were present in 114 cases (88%), down slightly from 2010. Prescription opioids were found in 62% of the cases, down sharply from 2010. Those most frequently mentioned in toxicology reports were: methadone (33%), essentially the same as 2010, followed by oxycodone (19%), and hydrocodone (15%), both down from 2010. (National studies suggest the methadone deaths result from methadone originally prescribed for pain, *not* methadone diverted from drug abuse treatment programs.) Heroin was present in 35% of the cases, an increase from 31% from 2010. There were 6 deaths where heroin was present in the absence of other central nervous system depressant drugs.

Sedatives, such as benzodiazepines, were the second most frequently mentioned drug in the toxicology reports, with about 75% of decedents having one or more in their systems at the time of death.

Cocaine was present in 41% of the cases, a sharp increase from 30% in 2010.

Alcohol was mentioned in 23% of the cases, the same as in 2010.

Nearly half of the decedents (49%) had *both* a benzodiazepine *and* a prescription opioid in their system, down slightly from 57% in 2010; 106 (82%) had two or more central nervous system depressants in their systems when they died. It is well-established that the concurrent use of drugs that depress the central nervous system can be extremely hazardous and result in death.

The PDR is conducted by the Center for Interventions, Treatment & Addictions Research under contract with Public Health – Dayton & Montgomery County (PHDMC) with support from the Ohio Department of Health (ODH). The PDR is part of the Preventing Unintentional Drug Poisoning Project, which is funded by PHDMC and the ODH with injury prevention block funds from the Centers for Disease Control (CDC).

ISONING DEATH REV					
Total Casas	lon 1 Dec 21 2011		120		2010 Ca
Total Cases	Jan. 1 - Dec. 31, 2011		130		
MOGRAPHICS			1	ı	
Characteristic	Data Source				
		Category	Freq	Percent	2010 Per
Age	Death Certificate	<15 years	0		
		15-24 years	7	5%	
		25-34 years	29	22%	
		35-44 years	34	26%	
		45-54 years	41	32%	
		55-64 years	18	14%	
		65-74 years	1	1%	
		75+ years	0	0%	
Gender	Death Certificate	Male	77	59%	
		Female	53	41%	
Race	Death Certificate	White	113		
		Black	17	13%	
		Other	0		
Hispanic	Death Certificate	Hispanic/Latino	0		
Education *	Death Certificate	<high school<="" td=""><td>36</td><td></td><td></td></high>	36		
		HS graduate	89		
		College graduate	3	-	
		Post-graduate	1	1%	
Marital Status*	Death Certificate	Single	50		
		Married	40		
		Divorced	32		
		Separated	4		
		Widowed	3	2%	
a attr	5 11 6 115	Ever in US Armed	_	401	
Military ALTH	Death Certificate	Forces	5	4%	
ALIN					
Characteristic	Data Source				
			Freq	Percent	2010 Per
	1				
Physical Disability/Illness	Case Synopsis and		96	74%	
Heart Disease	Postmortem Report		73	56%	
Mental Disability/Illness					
Merical Disability/ IIII1633			30	23%	

STORY OF SUBSTANCE ABUS	E				
					2010 Case
		Total Cases	130		13
Substance Abuse	Data Source				
			Freq	Percent	2010 Perce
Any history			107	82%	75
Alcohol			23	18%	13
Cocaine			22	17%	12
Marijuana	Case Synopsis and		2	2%	!
Heroin	Postmortem Report		34	26%	20
Prescription opioids	rostinortem Report		35	27%	2
Benzodiazepines			21	16%	1
Other Prescription					
Medications			7	5%	3
Over-the-Counter					
Medications			0	0%	
ATH INVESTIGATION					
Characteristic	Data Source	Category			
			Freq	Percent	2010 Perce
Location of death	Case Synopsis	Decedent's home	69		
	, ,	Relative's home	0	0%	
		Friend's home	21	16%	1
		Place of work	1	1%	(
		School	0	0%	
		Hospital	29	22%	
		Drug tx facility	0	0%	
		Jail/detention area	0	0%	
		Public area	2	2%	
		Other	8	6%	
911 called	Case Synopsis	Yes	127	98%	9
Person reporting death*		Coroner	0	0%	
	Case Synopsis	Hospital physician	30	23%	1
		Mortician	0	0%	
		EMS/Police	99	75%	8
Possible prevention by use of opioid antagonist?	Case Synopsis Postmortem Report				
C. Spiela antagonisti	Toxicology Report		18	14%	1
XICOLOGY REPORT					
					2010 Cas
		Total Cases	130		1

Characteristic	Data Source	Category	Freq	Percent	2010 Percent
		Alcohol	30	23%	23%
		Cocaine	53	41%	30%
		Methamphetamine	2	2%	1%
		Heroin	46	35%	31%
Prescription Opioids		Any	81	62%	74%
		Oxycodone	25	19%	23%
		Hydrocodone	19	15%	24%
		Methadone	43	33%	32%
		Fentanyl	8	6%	7%
		Tramadol	5		6%
		Hydromorphone	0		1%
		Morphine	10		9%
		Other	4	3%	4%
Anti-Depressants		Any	41	32%	
Opioids		Any	114	88%	92%
Sedatives (including	Toxicology Report				
benzodiazepines)		Any	96		
Benzodiazepines		Any	84	65%	70%
Any Prescription Opioid +					
Any Benzodiazepine			64	49%	57%
Two or more of the following					
CNS depressants: alcohol,					
heroin, prescription opioids,					
and/or sedatives			107	82%	85%
Heroin + any other CNS					
depressant			40	31%	27%
Heroin without any other					
CNS depressant			6	5%	4%
Other Prescription		Any	48	37%	39%
Over-The-Counter		Any	25	19%	20%
Varifiable Valid Bressrinties					
Verifiable Valid Prescription	Investigator's Report				
for Controlled Drugs in	congator o meport			370/	320/
Toxicology Report				37%	33%

^{*}Total may not equal 130 because of missing data