Wright State University

CORE Scholar

Chemistry Student Publications

Chemistry

6-1-2012

New Primers Reveal the Presence of a Duplicate Histone H3 in the Marine Turtle Leech *Ozobranchus branchiatus*

Triet Minh Truong

Wright State University - Main Campus

Audrey E. McGowin Ph.D.

Wright State University - Main Campus, audrey.mcgowin@wright.edu

Philip Lavretsky

Jeffrey L. Peters Ph.D.

Wright State University - Main Campus, jeffrey.peters@wright.edu

Follow this and additional works at: https://corescholar.libraries.wright.edu/chem_student

Part of the [Chemistry Commons](#)

Repository Citation

Truong, T. M., McGowin, A. E., Lavretsky, P., & Peters, J. L. (2012). New Primers Reveal the Presence of a Duplicate Histone H3 in the Marine Turtle Leech *Ozobranchus branchiatus*. .
https://corescholar.libraries.wright.edu/chem_student/6

This Presentation is brought to you for free and open access by the Chemistry at CORE Scholar. It has been accepted for inclusion in Chemistry Student Publications by an authorized administrator of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

New primers reveal the presence of a duplicate histone H3 in the marine turtle leech *Ozobranchus branchiatus*

Triet M. Truong and Audrey E. McGowin, Ph.D.
Department of Chemistry

Philip Lavretsky and Jeffrey L Peters, PhD
Department of Biological Sciences

Introduction

Marine leeches, specific to sea turtles, have been implicated as potential vector organisms in the spread of fibropapillomatosis (FP), a pandemic neoplastic disease with green turtles (*Chelonia mydas*) having the highest affliction rate. Polymerase chain reaction identified two independent, seemingly functional histone H3 loci for marine turtle leeches *Ozobranchus branchiatus* collected from *C. mydas* in Florida and Hawaii. Primers were developed to amplify each product separately. These novel markers will be useful in identifying ectoparasites in FP research, evaluating other histone variants, and chromatin dynamics regulation studies.

Methods

❖ Methodology for sampling and morphological identification of *O. branchiatus* are given in McGowin *et al.* 2011.

❖ Nuclear protein coding-gene histone H3 were amplified using primers given in Rousset *et al.* (2007). When multiple products were visible in the chromatograms for the histone H3 gene, new reverse primers synthesized by Invitrogen Corporation (Carlsbad, CA, USA) were employed to obtain better sequencing results (Lavretsky *et al.* 2011):

Reverse 1:H3R1 (5'-CCAACCAAGTACGCCTCA-3')

Reverse 2:H3R2 (5'-CCAACCAAGTAAGCCTCG-3')

H3R1 and H3R2 have an annealing temperature of 55°C and functionally compatible with the H3af forward primer given in Rousset *et al.* 2007.

❖ Each 25-μL reaction mixture containing template, GoTaq® Green Master Mix, forward and reverse primers, and ddH₂O were prepared according to the GoTaq® Green Promega protocol as provided (<http://www.promega.com>).

❖ The PCR thermal regime for amplification was 94°C for 7 minutes, followed by 45 cycles of 40 s at 94°C, 40 s at a specific annealing temperature (52-55 °C), 45 s at 72 °C, and then a final extension of 7 min at 72°C using a PCR thermocycler (Eppendorf Mastercylers).

❖ Purification of PCR Products was done using Agencourt® Ampure® XP Protocol 000387v001. However, after washing with 70% ethanol twice, PCR products were eluted using ddH₂O instead of elution buffer.

❖ Amplification products were sequenced in both directions using the Sanger Sequencing Method. Sequencing master mix included 4.5 μL ddH₂O, 1.75 μL Buffer (5X), 1 μL primer, and 1 μL BigDye per reaction and ran at the following thermocycler settings: 96°C for 1 minute, followed by 30 cycles of 10 s at 96°C, 5 s at 50°C, and 4 minutes at 60°C, and then a final hold temperature at 4°C. Buffer (5X) and BigDye provided by BigDye® Terminator v3.1 Cycle Sequencing Kit.

❖ Alignment analysis of genetic sequences was done using Sequencer™ 4.9 (Gene Codes, Inc.). A consensus maximum parsimony tree was generated using the Close-Neighbor-Interchange algorithm (Nei and Kumar 2000) in MEGA5 (Tamura *et al.* 2011).

References

Lavretsky P, Truong TM, McGowin AE, Balazs GH, Peters J. New primers reveal the presence of a duplicate histone H3 in the marine turtle leech *Ozobranchus branchiatus*. *Conservation Genetics Resources* (2012), 4, 487-490.

McGowin AE, Truong TM, Corbett AM, Bagley DA, Ehrhart LM, Bresette MJ, Weege ST, Clark D. (2011) Genetic barcoding of marine leeches (*Ozobranchus* spp.) from Florida sea turtles and their divergence in host specificity. *Molecular Ecology Resources*, 11, 271– 278.

Nei M, Kumar S. (2000) Molecular evolution and phylogenetics. Oxford University Press, New York.

Rousset, V., Pleijel, F., Rouse, G. W., Erseus, C., & Siddall, M. E. (2007). A molecular phylogeny of annelids. *Cladistics*, 23, 41-63.

Tamura K & Nei M (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, 10, 512-526.

Tamura K, Peterson D, Peterson N, Stecher G, Nei M, and Kumar S (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution* (online).

Table 1 Estimates of evolutionary divergence (Φ_{ST}) between histone H3 loci and <i>Ozobranchus branchiatus</i> populations			
Histone H3 Haplotype	Hawaii R1	Florida R1	Hawaii R2
Hawaii R1	—	—	—
Florida R1	0.004	—	—
Hawaii R2	0.161	0.172	—
Florida R2	0.171	0.182	0.005

Conducted in MEGA5 (Tamura *et al.* 2011) using the Tamura 3-parameter model (Tamura 1992) with a gamma distribution (shape parameter = 5) and a total of 12 nucleotide sequences (259 base pairs/ sequence). Ambiguous positions were removed for each sequence pair.

Images of *O. branchiatus* from each sampling site are displayed above. Except for the *O. branchiatus* associated with the Indian River Lagoon location (specimen was collected on the same *C. mydas* as those sequenced), all photographed leeches were sequenced in this study.

Table 2 Histone H3 sequence accession data for specimens analyzed.		
Higher taxa	Species	GenBank accession no.
ANNELIDA Clitellata	Haplotaxida	
	<i>Allolobophora chlorotica</i>	FJ214261 [†] , FJ374780 [†]
	<i>Antarctodrilus proboscidea</i>	DQ779716 [‡]
	<i>Aporrectodea caliginosa</i>	FJ214248 [†] , FJ374784 [†]
	<i>Aporrectodea icterica</i>	FJ214249 [†] , FJ374781 [†]
	<i>Aporrectodea longa</i>	FJ214250 [†] , FJ214251 [†]
	<i>Aporrectodea rosea</i>	FJ214252 [†] , FJ214262 [†]
	<i>Aporrectodea tuberculata</i>	FJ214268 [†] , FJ374783 [†]
	<i>Dendrobaena attemsi</i>	FJ214254 [†]
	<i>Dendrobaena clujensis</i>	FJ214253 [†]
	<i>Dendrobaena hortensis</i>	FJ214259 [†]
	<i>Dendrobaena veneta</i>	FJ214264 [†]
	<i>Dendrodrilus rubidus</i>	FJ214239 [†] , FJ374782 [†]
	<i>Eisenia andrei</i>	DQ779728 [‡]
	<i>Eisenia fetida</i>	FJ214246 [†] , FJ214247 [†] , FJ214258 [†]
	<i>Eisenia lucens</i>	FJ214255 [†]
	<i>Eisenoides carolinensis</i>	FJ214256 [†]
	<i>Helodrilus oculatus</i>	FJ214245 [†] , FJ214271 [†]
	<i>Heronidrilus gravidus</i>	DQ779736 [‡]
	<i>Lumbricus castaneus</i>	FJ214244 [†] , FJ214272 [†]
	<i>Lumbricus festivus</i>	FJ214243 [†] , FJ214270 [†]
	<i>Lumbricus rubellus</i>	FJ214242 [†] , FJ214257 [†]
	<i>Lumbricus terrestris</i>	AF185262 [§] , FJ214240 [†] , FJ214241 [†] , FJ214260 [†]
	<i>Octolasion lacteum</i>	FJ214267 [†]
	<i>Octolasion tyrtaeum</i>	FJ214263 [†] , FJ214265 [†] , FJ214266 [†] , FJ214269 [†]
Arihynchobdellida	<i>Erpobdella octoculata</i>	DQ779729 [‡]
	<i>Hirudo medicinalis</i>	DQ779738 [‡]
Rhynchobdellida	<i>Glossiphonia complanata</i>	DQ779733 [‡]
	<i>Helobdella stagnalis</i>	DQ779735 [‡]
	<i>Ozobranchus branchiatus</i> [*]	JQ070406 ^{‡123} , JQ070407 ^{‡45} , JQ070408 ^{‡4} , JQ070409 ^{‡1} , JQ070410 ^{‡23} , JQ070411 ^{‡4} , JQ070412 ^{‡5} , JQ0704013 ^{‡6}

[†]Lund et al. 2008; [§]Rouse et al. 1999; [‡]Rousset et al. 2007; ^{*}This study; histone H3R1[‡]; histone H3R2[‡]

Florida sites: Grassy Key MM 57[†]; Indian River Lagoon[‡]; Hutchinson Island[‡]

Hawaii sites: Lahaina, Maui[‡]; Kalapolepo Fish Pond, Maui[‡]; West Loch, Oahu[‡]

Figure 1 Maximum parsimony analysis of 59 annelid histone H3 sequences conducted in MEGA5 (Tamura *et al.* 2011) using 1000 bootstrap replicates. GenBank accession numbers along with the corresponding taxa names and associated authors provided in Table 1.

Florida sites: Grassy Key MM 57[†]; Indian River Lagoon[‡]; Hutchinson Island[‡]

Hawaii sites: Lahaina, Maui[‡]; Kalapolepo Fish Pond, Maui[‡]; West Loch, Oahu[‡]

Results and Conclusions

❖ It is possible not all loci composing the histone H3 complex were identified and that pooling individuals resulted in false-positives for heterozygotes, but sequence divergence estimates (Φ_{ST}) ranging from 0.161 to 0.182 (Table 1) suggested at least two divergent loci (histone H3R1 and H3R2).

❖ The two well supported ($\geq 99\%$ bootstrap support) monophyletic groups (exclusively including either H3R1/H3R2 haplotypes) confirms the new locus-specific primers amplified two divergent loci (Figure 1). The two loci are sister to one another (99% bootstrap support) rather than to other annelids, suggesting the duplication event occurred within the Ozobranchidae lineage.

❖ Despite being phylogenetically distinctive, genetic diversity based upon location showed low levels of differentiation between Hawaiian and Floridian samples, with Φ_{ST} estimates of 0.004 and 0.005 for histone H3R1 and H3R2, respectively (Table 1). Indeed, there were two fixed differences between alleles sampled from Florida and Hawaii at H3R1 (Figure 1).

Acknowledgements

This research was funded by the Women in Science Giving Circle (WISGC). Special thanks to George H. Balazs (NOAA's Pacific Islands Fisheries Science Center of the National Marine Fisheries Service, Honolulu) who collected all leeches from Hawaii. Ryan Butts (The Sea Turtle Hospital, Marathon) for samples from Grassy Key, Dean Bagley (University of Central Florida) for leeches from Indian River Lagoon, and Dave Clark and Stephen T. Weege (Inwater Research Group, Inc.) for leeches from the St. Lucie Nuclear Power Plant. John O. Stireman III, PhD (Wright State University) generously provided his microscope camera throughout the research.

<http://www.chm.wright.edu/mcgowin/research/index.html>