

1-2010


Traveling Wave Solutions for a Nonlocal Reaction-Diffusion Model of Influenza A Drift

Joaquin Riviera

Yi Li

Wright State University - Main Campus, yi.li@csun.edu

Follow this and additional works at: <https://corescholar.libraries.wright.edu/math>

 Part of the [Applied Mathematics Commons](#), [Applied Statistics Commons](#), and the [Mathematics Commons](#)

Repository Citation

Riviera, J., & Li, Y. (2010). Traveling Wave Solutions for a Nonlocal Reaction-Diffusion Model of Influenza A Drift. *Discrete and Continuous Dynamical Systems - Series B (DCDS-B)*, 13 (1), 157-174.
<https://corescholar.libraries.wright.edu/math/72>

This Article is brought to you for free and open access by the Mathematics and Statistics department at CORE Scholar. It has been accepted for inclusion in Mathematics and Statistics Faculty Publications by an authorized administrator of CORE Scholar. For more information, please contact corescholar@www.libraries.wright.edu, library-corescholar@wright.edu.

EXISTENCE OF TRAVELING WAVE SOLUTIONS FOR A NONLOCAL REACTION-DIFFUSION MODEL OF INFLUENZA A DRIFT

JOAQUIN RIVIERA

Department of Mathematics
Colgate University
Hamilton, NY 13346, USA

YI LI

Department of Mathematics
University of Iowa
Iowa City, IA 52240, USA

and
Department of Mathematics
Xian Jiaotong University
Xian, China

(Communicated by Yuan Lou)

ABSTRACT. In this paper we discuss the existence of traveling wave solutions for a nonlocal reaction-diffusion model of Influenza A proposed in Lin et. al. (2003). The proof for the existence of the traveling wave takes advantage of the different time scales between the evolution of the disease and the progress of the disease in the population. Under this framework we are able to use the techniques from geometric singular perturbation theory to prove the existence of the traveling wave.

1. Introduction. Recently work has been developed in ecological and epidemiological models using integro-differential equations to model spatial processes. Kendall [9], studied the traveling wave solutions for an integro-differential SIR model, where the integral term models the nonlocal contacts in among the susceptible and infectious populations. In his model Kendall assume that the rate of infection is given by:

$$\beta \int_{-\infty}^{\infty} I(x, t) K(x - y) dy, \quad (1)$$

where $I(x, t)$ represent the density of the infected individuals at time t , and $K(x)$ is a kernel function of mass one that measure the contribution of the infected individuals at location y to the infection of susceptible individuals at location x . The integral models the fact that two individuals can influence each other even if their separation in subtypes is large. For more about these epidemiological models the reader should

2000 *Mathematics Subject Classification.* Primary: 58F15, 58F17; Secondary: 53C35.

Key words and phrases. Dimension theory, Poincaré recurrences, multifractal analysis.

The first author is supported by a grant from the Sloan Foundation and the Graduate College of the University of Iowa.

consult [14, 16] for a discussion about the use of integro-differential equations in the modeling of spreading diseases.

The recorded patterns of influenza A infection contain two important phenomena, the first being the almost identical annual epidemics which occur in most countries, and the second is the extensive pandemics which occurs approximately every 25-30 years. The Influenza A pandemics are mostly caused by appearance of new subtypes of the virus. The new variants of the virus are the result of changes in the amino acids of the HA and NA proteins between pandemics [15]. This process is known as *antigenic drift*. The approach followed in [13] to model antigenic drift was by allowing the virus to diffuse along a one-dimensional axis of variants x .

In this situation one natural question is the behavior of solution $u(x, t)$ as $t \rightarrow +\infty$. In particular, we are interested to know about the existence of traveling wave solutions of the form $u(\xi) = u(x - ct)$ such that $u \rightarrow 0$ as $\xi \rightarrow \pm\infty$. Here $c > 0$ is referred to as the speed of the pulse. The study of the existence, and stability of traveling waves have generated a vast amount literature on the subject. We refer to the books by Volpert et al [18] and Fife [5] for a more general discussion about traveling waves.

Many authors have studied the existence and stability of traveling fronts in reaction-diffusion equations [2, 4, 7, 10, 11]. In [1, 2] a systematic study was carried out for the existence of traveling wave solutions for a semilinear parabolic equation in higher dimensional cylinders. In [6, 12] the authors expanded the local stability results for reaction-diffusion equations in various weighted Banach spaces.

In this paper we discuss the existence of traveling wave solutions for an integro-differential model for the spread of Influenza A proposed in [13]. After some rescaling the model is given by the following system of equations:

$$\frac{dS}{dt} = e(1 - S) - R_0SI, \quad (2a)$$

$$\frac{\partial i}{\partial t} = \epsilon \frac{\partial^2 i}{\partial x^2} + R_0i \int_{-\infty}^x K(x-y)r(y,t)dy + R_0Si - i, \quad (2b)$$

$$\frac{\partial r}{\partial t} = -rR_0 \int_x^{\infty} K(x-y)i(y,t)dy + (1-e)i - er, \quad (2c)$$

The variables in the model are represented by: $S(t)$ represent the number of susceptible host at time t , while $i(x, t)$ and $r(x, t)$ are the distribution of infected individuals with variant x and recovered individuals from variant x at time t , respectively. Observe that the nonlinear terms in (2b,2c) are given by the nonlocal terms. The nonlocal term in (2b) represent the per capita growth rate of infection by hosts carrying variant x , and the diffusion term in (2b) is the drift of the virus. In [13], the authors carried out a preliminary analysis that suggested the existence of traveling wave solutions for the system. In our paper, we want to formalize those findings in a more mathematical way, at least for certain class of kernels. In [13], the authors used the Monoid kernel $K(v) = v/(v+a)$, while in this paper we proposed a more standard exponential kernel that satisfied the conditions on [13].

The proof for the existence of traveling waves solutions for (2) uses geometrical singular perturbation theory [3, 8] to study the dynamics of the system. The analysis of the proof for the existence of the traveling wave solutions takes the advantage of the smallness of the diffusion coefficient. It is realistic to assume that the kernel, because recovered individuals are protected against close by variants of the virus

than from those further away has the form:

$$K(x) = 1 - e^{-ax}, x > 0 \quad (3)$$

for some $a > 0$. The parameter a measures the nonlocality of the disease transmission, so that for a large this kernel gives a strong nonlocality, i.e. if a is large strains of the value that are far away in the variants spaces become more important. The main result of this paper is the existence of traveling wave solution for (2). The first part of the argument considers the extended system, that is (2) and the equations given by the new variables (5) below. Thus first, by studying the slow-fast dynamics of the system for the case when $\epsilon = 0$ we establish the existence of the traveling wave in the critical manifold.

Theorem 1.1. *The unperturbed system when $\epsilon = 0$ has a heteroclinic solution connecting the quasi steady-states $(0, 0, 0, \mathcal{I}_{-\infty})$ and $(0, 0, \mathcal{R}_{+\infty}, 0)$ where $\mathcal{R}_{+\infty} = \frac{2(1-2e)}{cR_0}$, $\mathcal{I}_{-\infty} = \frac{2(1-2e)}{cR_0}$, and $S = \frac{2e}{R_0}$.*

For the perturbed system (2), we construct the so-called slow manifold in an ϵ -neighborhood of the critical manifold. By a careful study of the asymptotic behavior of the solution we were able to transform the problem of the existence of the solution to a corresponding fixed point problem of a nonlinear map. We apply the Contraction Mapping Principle in some weighted Banach space to show that the existence of the traveling wave solution is also true for $\epsilon > 0$, but small.

Theorem 1.2. *System (51) has a traveling wave solution, and furthermore the solution is in a neighborhood of the unperturbed solution.*

The rest of the paper is organized in the following manner: in Section 2 we discuss the structure of the traveling wave, and some preliminaries analysis for the existence of the traveling wave solutions for the whole system. The construction of the traveling wave solutions on the critical manifold. In Section 3 we prove the robustness of the traveling wave for the perturbed system.

2. Structure of the traveling waves. In traveling wave form, with $z = x - ct$, the original system can be reformulated as:

$$0 = e(1 - S^*) - R_0 S^* I, \quad (4a)$$

$$-c \frac{di}{dz} = \epsilon \frac{d^2 i}{dz^2} + R_0 i \int_{-\infty}^z K(z-w)r(w)dw + R_0 S^* i - i, \quad (4b)$$

$$-c \frac{dr}{dz} = -rR_0 \int_z^{\infty} K(w-z)i(w)dw + (1-e)i - er, \quad (4c)$$

with the boundary conditions $(i(\pm\infty), r(\pm\infty)) = (0, 0)$, such traveling wave solutions are sometime referred to as traveling pulses. Recall that we are interested in finding a homoclinic orbit connecting the origin at $z \rightarrow \pm\infty$. We assume that, in this framework, S^* represents a quasi steady-state for the susceptible population, and we only need to focus on the equations (4b) and (4c). Defining the new variables:

$$R(z) = \int_{-\infty}^z K(z-w)r(w)dw, \quad (5a)$$

$$I(z) = \int_z^{\infty} K(w-z)i(w)dw. \quad (5b)$$

Observe that $K'(x) = a(1 - K(x))$, thus by computing the derivatives of R and I we obtain:

$$\begin{aligned} R'(z) &= \int_{-\infty}^z K'(z-w)r(w)dw \\ &= \int_{-\infty}^z a(1 - K(z-w))r(w)dw \\ &= a \int_{-\infty}^z r(w)dw - a \int_{-\infty}^z K(z-w)r(w)dw \end{aligned}$$

Thus, by computing the second derivative we have that $R(u)$ satisfies the following second order ODE:

$$R''(z) = ar(z) - aR'(z). \quad (6)$$

Similarly, we obtain that $I(u)$ satisfies the second ODE:

$$I''(z) = ai(z) + aI'(z). \quad (7)$$

Adding these new equations to (4b-c), and replacing the integral terms by the new variables (5) we obtain the following system of equations.

$$-c \frac{di}{dz} = \epsilon \frac{d^2i}{dz^2} + R_0Ri + R_0S^*i - i, \quad (8a)$$

$$-c \frac{dr}{dz} = -rR_0I + (1 - e)i - er, \quad (8b)$$

$$\frac{d^2R}{dz^2} = ar - aR', \quad (8c)$$

$$\frac{d^2I}{dz^2} = ai + aI'. \quad (8d)$$

The trade off of doing this is that the number of equations is increased from two to four. Analytical experiments suggest that ϵ is small while a is large when compare to ϵ . Thus, we assume that the cross-protection parameter a is inversely proportional to the mutation rate ϵ , that is $a = \frac{m}{\epsilon}$. Converting system (8) to a first order system of equations, we have

$$\epsilon i_1' = -ci_1 + (1 - R_0S^*)i - R_0Ri, \quad (9a)$$

$$\epsilon R_1' = m(r - R_1), \quad (9b)$$

$$\epsilon I_1' = m(i + I_1), \quad (9c)$$

$$i' = i_1, \quad (9d)$$

$$r' = \frac{R_0rI + (e - 1)i + er}{c}, \quad (9e)$$

$$R' = R_1, \quad (9f)$$

$$I' = I_1. \quad (9g)$$

Note that the problem of finding a homoclinic orbit for the original system (2) or (4) connecting the origin at $\pm\infty$, becomes a problem of finding a heteroclinic orbit for the system (9) that connects the steady-states solutions $(0, 0, 0, 0, 0, 0, I_{-\infty})$ to $(0, 0, 0, 0, 0, R_{+\infty}, 0)$, for some $I_{-\infty} > 0$, and $R_{+\infty} > 0$

2.1. Preliminaries. First we want to show that $1 - R_0S^* > 0$. Suppose on the contrary that $1 - R_0S^* \leq 0$. Integrating (8a) from $-\infty$ to z we obtain

$$\epsilon i'|_{-\infty}^z + ci|_{-\infty}^z + - \int_{-\infty}^z (-1 + R_0S^* + R_0R)i = 0 \quad (10)$$

So,

$$\epsilon i' + ci + \int_{-\infty}^z (-1 + R_0S^* + R_0R)i = 0 \quad (11)$$

Thus, $-1 + R_0S^* + R_0R \geq 0$ since $1 - R_0S^* \leq 0$, so we have that

$$\epsilon i' + ci \leq 0 \quad (12)$$

or

$$(e^{\frac{c}{\epsilon}z}i)' \leq 0 \quad (13)$$

Thus, integrating (13) from z to 0,

$$e^{\frac{c}{\epsilon}z}i(z) \geq i(0) \quad (14)$$

or

$$i(z) \geq i(0)e^{-\frac{c}{\epsilon}z} \quad (15)$$

But $e^{-\frac{c}{\epsilon}z} \rightarrow +\infty$ as $z \rightarrow -\infty$, which is a contradiction. Hence we must have $1 - R_0S^* > 0$.

To help constructing the weighted Banach spaces in Section 3 let us compute the eigenvalues of the asymptotic systems at $(0, 0, 0, 0, 0, 0, \mathcal{I}_{-\infty})$ and at $(0, 0, 0, 0, 0, \mathcal{R}_{+\infty}, 0)$. At $-\infty$ the linear system is given by:

$$\begin{bmatrix} i_1 \\ R_1 \\ I_1 \\ i \\ r \\ R \\ I \end{bmatrix}' = \begin{bmatrix} -\frac{c}{\epsilon} & 0 & 0 & \frac{1-R_0S^*}{\epsilon} & 0 & 0 & 0 \\ 0 & -\frac{m}{\epsilon} & 0 & 0 & \frac{m}{\epsilon} & 0 & 0 \\ 0 & 0 & \frac{m}{\epsilon} & \frac{m}{\epsilon} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{e-1}{c} & \frac{R_0\mathcal{I}_{-\infty}+e}{c} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} i_1 \\ R_1 \\ I_1 \\ i \\ r \\ R \\ I \end{bmatrix} = A_{-\infty} \begin{bmatrix} i_1 \\ R_1 \\ I_1 \\ i \\ r \\ R \\ I \end{bmatrix} \quad (16)$$

And the set of eigenvalues for the matrix $A_{-\infty}$ is given by,

$$\sigma_{-\infty} = \left\{ 0, 0, \frac{m}{\epsilon}, -\frac{c \pm \sqrt{c^2 + 4\epsilon - 4\epsilon R_0S^*}}{2\epsilon}, -\frac{m}{\epsilon}, \frac{R_0\mathcal{I}_{-\infty} + e}{c} \right\}. \quad (17)$$

Observe that there are two negative eigenvalues, three positive eigenvalues, and two zero eigenvalues. This implies that at $(0, 0, 0, 0, 0, 0, \mathcal{I}_{-\infty})$ there is a 3-dimensional unstable manifold, a 2-dimensional stable manifold, and there is also a 2-dimensional center manifold.

Similarly, for the asymptotic system at $+\infty$ the linearized matrix is given by

$$\mathcal{A}_{+\infty} = \begin{bmatrix} -\frac{c}{\epsilon} & 0 & 0 & \frac{1-R_0S^*-R_0\mathcal{R}_{+\infty}}{\epsilon} & 0 & 0 & 0 \\ 0 & -\frac{m}{\epsilon} & 0 & 0 & \frac{m}{\epsilon} & 0 & 0 \\ 0 & 0 & \frac{m}{\epsilon} & \frac{m}{\epsilon} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{e-1}{c} & \frac{e}{c} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The eigenvalues of $\mathcal{A}_{+\infty}$ are given in the following set

$$\sigma_{+\infty} = \left\{ 0, 0, \frac{m}{\epsilon}, -\frac{c \pm \sqrt{c^2 + 4\epsilon - 4\epsilon R_0 S^* - 4\epsilon R_0 \mathcal{R}_{+\infty}}}{2\epsilon}, -\frac{m}{\epsilon}, \frac{e}{c} \right\}. \quad (18)$$

Thus, there are three positive eigenvalues, two negatives and two zero eigenvalues. From (17), and (18) we obtain that:

$$-\frac{c - \sqrt{c^2 + 4\epsilon - 4\epsilon R_0 S^*}}{2\epsilon} \quad (19)$$

and

$$-\frac{c + \sqrt{c^2 + 4\epsilon - 4\epsilon R_0 S^* - 4\epsilon R_0 \mathcal{R}_{+\infty}}}{2\epsilon} \quad (20)$$

the two eigenvalues that we will use to estimate the weights in Section 3.

Remark 1. We can think of the heteroclinic orbit connecting $(0, 0, 0, 0, 0, 0, I_{-\infty})$ and $(0, 0, 0, 0, 0, R_{+\infty}, 0)$ as the intersection of the unstable manifold of the asymptotic system at $-\infty$ with stable manifold of the asymptotic system at $+\infty$. Observe that the critical speed of the wave $c = c(\epsilon)$ depends on the parameter ϵ , as it is illustrated in (19), and (20)

2.2. Construction of the traveling wave solution. Note that the system (9) can be formulated in slow-fast variables by setting the new variables $x = (i_1, R_1, I_1)^T$ and $y = (i, r, R, I)^T$, which are the fast and slow variable, respectively. Thus, rewriting system (9) as:

$$\epsilon x' = f(x, y, \epsilon), \quad (21)$$

$$y' = g(x, y, \epsilon), \quad (22)$$

where the functions f , and g are given by the right hand sides of system (9). The motivation for this separation is to take advantage of the geometric singular perturbation methods to study the dynamics for the fast and slow system.

Setting $\epsilon = 0$ we take the critical manifold to be the set given by:

$$\mathcal{M}_0 = \left\{ (i, r, R, I, i_1, R_1, I_1) : i_1 = \frac{i(1 - R_0 S^* - R_0 R)}{c}, R_1 = r, I_1 = -i \right\}. \quad (23)$$

for some $c > 0$. Thus, we obtain the equations for the slow flow on the critical manifold \mathcal{M}_0 by substituting the relations on \mathcal{M}_0 into the remaining equations:

$$i' = \frac{i(1 - R_0 S^* - R_0 R)}{c}, \quad (24a)$$

$$r' = \frac{R_0 r I + (e - 1)i + er}{c}, \quad (24b)$$

$$R' = r, \quad (24c)$$

$$I' = -i, \quad (24d)$$

The only condition that we need check on \mathcal{M}_0 is that it is normally hyperbolic.

Lemma 2.1. *The critical Manifold \mathcal{M}_0 is normally hyperbolic relative to the fast system.*

Proof. Note that we only need to show that the linearization for the fast system evaluated on \mathcal{M}_0 has four eigenvalues with zero real part and three with nonzero

real part. Thus, it suffices to show that $D_x f(\hat{x}, \hat{y}, 0)$ has three eigenvalues with $Re(\lambda) \neq 0$ for any $(\hat{x}, \hat{y}) \in \mathcal{M}_0$. The Jacobian matrix for $f(\hat{x}, \hat{y}, 0)$ is given by:

$$J = \begin{bmatrix} -c & 0 & 0 \\ 0 & -m & 0 \\ 0 & 0 & m \end{bmatrix},$$

which has nonzero eigenvalues $\lambda = -c, \pm m$. Hence \mathcal{M}_0 is normally hyperbolic. \square

From the Lemma we have that the critical manifold has a 2 dimensional stable manifold and a 1 dimensional unstable manifold. Thus by applying all of the Fenichel's Theorem there is, for some $\epsilon > 0$ but small a manifold \mathcal{M}_ϵ which is diffeomorphic and close to \mathcal{M}_0 . Before we deal with the perturbed system in \mathcal{M}_ϵ , we will discuss the dynamics on the critical manifold, and then we will connect it to the dynamics on the slow manifold.

3. Dynamics of the model.

3.1. Dynamics on the critical manifold. In this section we show the existence of a traveling wave solution of (24) connecting the equilibrium points $(0, 0, 0, \mathcal{I}_{-\infty})$ and $(0, 0, \mathcal{R}^{+\infty}, 0)$. Note that this is a four dimensional system of equations, so the typical techniques to establish the existence of heteroclinic curves, like phase-plane analysis, are not applicable here. In order to tackle the problem we will rewrite the system as a single equation of fourth order, then construct a solution for that equation.

Note that by substituting the equations $I' = -i$, and $R' = r$ into the equations for i' and r' we obtain the new second order equations:

$$I'' = \frac{I'(1 - R_0 S^* - R_0 R)}{c}, \quad (25)$$

$$R'' = \frac{-I(e - 1) + eR' + R_0 I R'}{c}. \quad (26)$$

Now, observe that we can solve for R in the first equation obtaining the relation:

$$R = \frac{I'(1 - R_0 S^*) - cI''}{R_0 I'}, \quad (27)$$

thus, by computing the first and second order derivative of R , by using (27), i.e.

$$R' = \left[\frac{(I''(1 - R_0 S^*) - cI''')R_0 I' - R_0 I''(I'(1 - R_0 S^*) - cI'')}{R_0^2 I''^2} \right], \quad (28)$$

and

$$R'' = \frac{-cI^{(4)}I' + (1 - R_0 S)R_0 I' + cR_0 I''}{R_0^4 I''^2}. \quad (29)$$

Thus by substituting this into the second equation we have the following fourth order differential equations on $I(z)$:

$$I^{(4)} - I^{(3)} \left(\frac{3I^{(2)}}{I^{(1)}} + \frac{e + R_0 I}{c} \right) + 2 \frac{I^{(2)^3}}{I^{(1)^2}} + \frac{(e + R_0 I)I^{(2)^2}}{cI^{(1)}} - \frac{R_0(e - 1)I^{(1)^2}}{c^2} = 0, \quad (30)$$

where $I^{(j)}$ denotes the j -th derivative of I with respect to z . Observe that now we only need to find a heteroclinic solution of (30) that connects the points 0 to \mathcal{I}^* , where $\mathcal{I}^* = \mathcal{I}_{-\infty}$. This may seem like a more difficult problem than the original system (24), since it is of higher order nonlinear equation. But by a change of variables we are able to reduce the equation (30) to a second order singular equation.

Let $v = I(z)$ and $\frac{dI}{dz} = g(v)$. Computing all the derivatives in (30), and combining all the terms we have that $g(v)$ is the solution of the following singular second order ordinary differential equation:

$$g(v) \frac{d^2 g(v)}{dv^2} - \left(\frac{R_0 v + e}{c} \right) \frac{dg(v)}{dv} + \frac{R_0 g(v)}{c} - \frac{R_0(e-1)}{c^2} v + C_2 = 0 \quad (31)$$

on $[0, \mathcal{I}^*]$, where $z = \int \frac{1}{g(v)} dv + C_1$, for some constants C_1 and C_2 . Notice that singular nature of equation is not due to limiting behavior as $\epsilon \rightarrow 0$, but to the fact the equation changes its order at any point $v_0 \in (0, \mathcal{I}^*]$ such that $g(v_0) = 0$.

By using the asymptotic behavior of $I(z)$ at $\pm\infty$ we can obtain the boundary conditions for (31) in $[0, \mathcal{I}^*]$. Thus, as $z \rightarrow -\infty$, it follows that $v \rightarrow \mathcal{I}^*$, so $g(v) \rightarrow g(\mathcal{I}^*) = 0$. Similarly, as $z \rightarrow +\infty$ we have that $g(v) \rightarrow 0$. Thus, the boundary conditions for (31) are $g(0) = 0 = g(\mathcal{I}^*)$.

From equation (24d) we obtain that $I(z)$ is a decreasing function and recall that $g(v) = \frac{dI}{dz} < 0$, and since $g(0) = 0 = g(\mathcal{I}^*)$ it must be the case that there is a v_0 so that $\min_{v \in [0, \mathcal{I}^*]} g(v) = g(v_0)$. By (31) we can obtain the following approximations for the function near 0 and \mathcal{I}^* :

$$g(v) = -Mv^\alpha, \quad (32a)$$

$$g(v) = -N(\mathcal{I}^* - v)^\beta, \quad (32b)$$

where M , and N are positive constants, and the exponents α , and β are to be determined.

We now substitute (32a) and (32b) into (31) to estimate the exponents α , and β . Thus, for (32a) we obtain:

$$(\alpha - 1)\alpha M^2 v^{2\alpha-2} + \frac{e}{c}\alpha M v^{\alpha-1} - \frac{R_0}{c} M v^\alpha - \frac{R_0(e-1)}{c} v + C_2 + h.o.t = 0, \quad (33)$$

Multiplying (33) by $v^{2-2\alpha}$ we obtain,

$$\begin{aligned} (\alpha - 1)\alpha M^2 + \frac{e}{c}\alpha M v^{1-\alpha} - \frac{R_0}{c} M v^{2-\alpha} \\ - \frac{R_0(e-1)}{c} v^{3-2\alpha} + C_2 v^{2-2\alpha} + h.o.t = 0, \end{aligned} \quad (34)$$

where $h.o.t$ denotes the remaining higher order terms. So, if $\alpha > 1$ we have that

$$C_2 = 0, \frac{e}{c}M = 0 \Rightarrow M = 0. \quad (35)$$

Now, if $\alpha < 1$ we have from (34) that

$$M^2 = 0 \Rightarrow M = 0. \quad (36)$$

Therefore, $\alpha = 1$. Similarly, we can deduce that $\beta = 1$.

Considering the test solution:

$$g(v) = -Mv(\mathcal{I}^* - v), \quad (37)$$

for some $M > 0$. Substituting (37) into (31) and solving for the constants terms C_2 , M , and \mathcal{I}^* to get a solution. Thus, we obtain the constants:

$$C_2 = \frac{e(2e-1)}{c^2}, \quad (38a)$$

$$M = \frac{R_0}{2c}, \quad (38b)$$

$$\mathcal{I}^* = \frac{(1-2e)}{R_0}. \quad (38c)$$

Considering the function $g(v) = -Mv(\mathcal{I}^* - v)$ as a test solution, where M , and \mathcal{I}^* are given by (38). By using this function and by reversing the change of variables we obtain the following:

$$z = -\frac{1}{Mf} \ln \left(\frac{f}{\mathcal{I}^* - f} \right), \quad (39)$$

for some choice of C_1 , and $f = I(z)$. Thus after solving (39) for f we obtain that

$$f = I(z) = \frac{e^{-zMT^*} \mathcal{I}^*}{e^{-zMT^*} + 1}. \quad (40)$$

Thus substituting the constants (38) into (40), and letting $\rho = \frac{1-2e}{c}$, we can simplify (40) as

$$I(z) = \frac{2c\rho}{R_0(e^{\rho z} + 1)}. \quad (41)$$

From (41) we can obtain $R(z)$ just by substituting the first and second order derivative of (41) into (27), so

$$R(z) = -\frac{\rho c e^{\rho z} + \rho c - e^{\rho z} - 1 + R_0 S e^{\rho z} + R_0 S^*}{(e^{\rho z} + 1)R_0}. \quad (42)$$

Since we are looking for such a solution $R(z)$ with the property $\lim_{z \rightarrow -\infty} R(z) = 0$, we must have $\rho c - 1 + R_0 S^* = 0$. And by solving for S^* we obtain $S^* = \frac{2e}{R_0}$. After simplifying (42) we have

$$R(z) = \frac{2\rho c e^{\rho z}}{(e^{\rho z} + 1)R_0}. \quad (43)$$

Finally from (24c) and (24d) we can get the functions $r(z)$ and $i(z)$

$$r(z) = \frac{2\rho^2 c e^{\rho z}}{(e^{\rho z} + 1)^2 R_0}, \quad (44a)$$

$$i(z) = \frac{2\rho^2 c e^{\rho z}}{(e^{\rho z} + 1)^2 R_0}. \quad (44b)$$

It is easy to check that the set $\{i(z), r(z), R(z), I(z)\}$ are solutions of the system (24) with the correct asymptotics. Hence, we have just constructed heteroclinic orbit on the critical manifold \mathcal{M}_0 that connects the points $(0, 0, 0, \mathcal{I}_{-\infty})$ and $(0, 0, \mathcal{R}_{+\infty}, 0)$.

The fast system is given by

$$i'_1 = -ci_1 + (1 - R_0S^*)i - R_0Ri \quad (45a)$$

$$R'_1 = m(r - R_1) \quad (45b)$$

$$I'_1 = m(i + I_1) \quad (45c)$$

$$i' = \epsilon i_1 \quad (45d)$$

$$r' = \epsilon \frac{R_0rI + (e - 1)i + er}{c} \quad (45e)$$

$$R' = \epsilon R_1 \quad (45f)$$

$$I' = \epsilon I_1 \quad (45g)$$

Note that if we let $\epsilon = 0$ the new reduced system is given by:

$$i'_1 = -ci_1 + (1 - R_0S^*)i - R_0Ri \quad (46a)$$

$$R'_1 = m(r - R_1) \quad (46b)$$

$$I'_1 = m(i + I_1) \quad (46c)$$

with (i, r, R, I) all constants. Thus, we can argue that the only dynamics of interest for the case when $\epsilon = 0$ is given by the slow system. For that reason the problem of interest when $\epsilon > 0$, but small, is to show that the traveling wave solution that we constructed in the previous sections persists for the perturbed problem.

3.2. Dynamics on the slow manifold. The purpose of this section is to study the dynamics of the perturbed system (8) for $\epsilon > 0$, but sufficiently small. By Lemma 2.1 we have that the critical manifold \mathcal{M}_0 is normally hyperbolic, so the existence of a slow manifold \mathcal{M}_ϵ follows by the invariant manifold theorem. The following characterization gives the relation between the slow manifold and the critical manifold:

$$i_1 = \frac{i(1 - R_0S^* - R_0R)}{c} + f_1(i, r, I, R, \epsilon) \quad (47a)$$

$$R_1 = r + f_2(i, r, I, R, \epsilon) \quad (47b)$$

$$I_1 = -i + f_3(i, r, I, R, \epsilon) \quad (47c)$$

Note that by invariant manifold theorem we have that the functions f_i are smooth. Thus, since ϵ is sufficiently small, and by taking the Taylor expansion of the f_i 's about $\epsilon = 0$, we can rewrite the perturbations terms as:

$$f_i(i, r, I, R, \epsilon) = f_{i_1}(i, r, I, R, 0) + \epsilon f_{i_2}(i, r, I, R, 0) + O(\epsilon^2), \quad (48)$$

for $i = 1, 2, 3$. Notice that $f_{i_1}(i, r, I, R, 0) = 0$. We also disregard the higher order terms for the moment. Then, we approximate the perturbed terms in (47) by only the linear terms in (48). Thus, the first order approximation for the perturbation is:

$$i_1 = \frac{i(1 - R_0S^* - R_0R)}{c} + \epsilon f_1(i, r, I, R), \quad (49a)$$

$$R_1 = r + \epsilon f_2(i, r, I, R), \quad (49b)$$

$$I_1 = -i + \epsilon f_3(i, r, I, R). \quad (49c)$$

The goal is to use this perturbations to study the flow on \mathcal{M}_ϵ . Now, we want to find explicit expression for the f_i on (49).

Substituting the perturbed terms (49) into (9a) and letting $\epsilon = 0$ we obtain the functional expressions for f_i . Thus, the first order perturbations are given by

$$i_1 = \frac{i(1 - R_0 S^* - R_0 R)}{c} + \epsilon \frac{i(1 + R_0 S^*(2R_0 R - R_0 S^* - 2) + R_0(R_0 R^2 - 2R - cr))}{-c^3}, \quad (50a)$$

$$R_1 = r + \epsilon \frac{i + -r + \rho c(i + r) - 2R_0 I r}{2mc}, \quad (50b)$$

$$I_1 = -i + \epsilon \frac{i(R_0 S^* + R_0 R - 1)}{mc}, \quad (50c)$$

and the flow is given by

$$i' = \frac{i(\rho c - R_0 R)}{c} + \epsilon \frac{i}{c^3}(1 - 2\rho c - R_0^2 R^2 + (1 - \rho c)(2R_0 R - 1 + \rho c) + 2R_0 R + cR_0 r), \quad (51a)$$

$$r' = \frac{1}{c}(R_0 I r + \frac{1}{2}(1 - \rho c)(i + r) - i), \quad (51b)$$

$$I' = -i + \epsilon \frac{R_0 R - \rho c}{mc}, \quad (51c)$$

$$R' = r + \epsilon \frac{i + \rho c i - r + \rho c r - 2R_0 I r}{2mc}, \quad (51d)$$

Note that system (51) is a regular perturbation of the slow system (24).

3.3. Construction of the singular solution. Now, we will look for solutions of (51) near the unperturbed solutions (41), (43), and (44). We will also show that the solutions on the slow manifold \mathcal{M}_ϵ are perturbations of the solutions of the unperturbed problem. In order to simplify the expressions we let $\epsilon = \frac{\delta c}{\rho}$, where $\delta \ll 1$. Thus, we obtain the following system of equations:

$$\frac{di}{du} = \frac{\delta^* R_0}{c\rho} r i + \frac{(\delta - \delta^*) R_0}{c\rho} r i + \frac{R_0(2\delta - 1)}{c} R i - \frac{\delta R_0^2}{c^2 \rho} R^2 i - \rho(\delta - 1)i, \quad (52a)$$

$$\frac{dr}{du} = \frac{R_0}{c} I r + \frac{1 - \rho c}{2c} r - \frac{1 + \rho c}{2c} i, \quad (52b)$$

$$\frac{dI}{du} = \frac{\delta R_0}{\rho m} R i - \frac{m + \delta c}{m} i, \quad (52c)$$

$$\frac{dR}{du} = \frac{\delta(1 + \rho c)}{2\rho m} i + \frac{\delta \rho c - 2\delta R_0 I - \delta}{2\rho m} r + r, \quad (52d)$$

where $0 < \delta^*$ is fixed.

First, we consider a linear system associated to (52)

$$\frac{di}{du} = \frac{\delta^* R_0}{c\rho} \tilde{r} \tilde{i} + \left[\rho(1 - \delta) + \frac{(\delta - \delta^*) R_0}{c\rho} r + \frac{R_0(2\delta - 1)}{c} \tilde{R} - \frac{\delta R_0^2}{c^2 \rho} \tilde{R}^2 \right] i, \quad (53a)$$

$$\frac{dr}{du} = \frac{R_0}{c} \tilde{I} r + \frac{1 - \rho c}{2c} r - \frac{1 + \rho c}{2c} i, \quad (53b)$$

$$\frac{dI}{du} = \frac{\delta R_0}{\rho m} \tilde{R} i - \frac{m + \delta c}{m} i, \quad (53c)$$

$$\frac{dR}{du} = \frac{\delta(1 + \rho c)}{2\rho m} i + \frac{\delta \rho c - 2\delta R_0 \tilde{I} - \delta}{2\rho m} r + r, \quad (53d)$$

where the functions $(\tilde{i}(u), \tilde{r}(u), \tilde{R}(u), \tilde{I}(u))$ represent known functions near the unperturbed solutions. Notice that by integrating the equations in (53) we obtain the following integral operator:

$$Tx = \begin{cases} \frac{\delta^* R_0}{c\rho} e^{\left(\int_{t_0}^u \alpha(s) ds\right)} \int_{-\infty}^u e^{\left(-\int_{t_0}^t \alpha(s) ds\right)} \tilde{i}(t) \tilde{r}(t) dt \\ \frac{1 + \rho c}{2c} \int_t^{+\infty} e^{-\frac{1}{2c} \int_t^{t_1} \beta(s) ds} i(t) dt \\ \int_u^{+\infty} \left(\frac{m + \delta c}{m} - \frac{\delta R_0 \tilde{R}(s)}{\rho m} \right) i(s) ds \\ \int_{-\infty}^u \frac{\delta(1 + \rho c)}{2\rho m} i(s) + \left[\frac{2\rho m - \delta + c\rho\delta}{2\rho m} - \frac{\delta R_0 \tilde{I}(s)}{\rho m} \right] r(s) ds \end{cases} \quad (54)$$

where $\alpha(t) = (1 - \delta)\rho + \frac{(\delta - \delta^*) R_0}{c\rho} r(t) - \frac{R_0}{c} (1 - 2\delta) R(t) - \frac{\delta R_0^2}{c^2 \rho} R(t)^2$, $\beta(t) = 2R_0 \tilde{I}(t) + 1 - \rho c$, and $x = (i, r, I, R)'$.

Let X be the space of continuous functions $C(\mathbb{R})$. Define the weighted Banach space to be the space X equipped with the following norm:

$$\|x\| = \sup_{u \in \mathbb{R}} \{(e^{\gamma_1 u} + e^{\gamma_2 u}) |x(u)|\} \quad (55)$$

where the exponents $\gamma_1 < 0$ and $\gamma_2 > 0$ are the asymptotic values of the growth function $\alpha(t)$ at $-\infty$, and $+\infty$, respectively. We denote the Banach space $(X, \|\cdot\|)$ by \mathbb{X} .

Let x_c be the unperturbed solution. If for $x \in \mathbb{X}^4$ such that $\|x - x_c\| \leq \delta$ for any $\delta \leq \delta^*$, then the asymptotic behavior of the function $\alpha(t)$ is given by

$$\alpha(t) = \begin{cases} -\rho(1 - 3\delta) + H_1(t), & \text{for } t \rightarrow +\infty \\ \rho(1 - \delta) + H_2(t), & \text{for } t \rightarrow -\infty \end{cases} \quad (56)$$

where the terms $H_i(t)$ decay exponentially fast as $t \rightarrow \pm\infty$.

Let $\mathcal{B}_{\delta_0} = \{(i, r, I, R) \in \mathbb{X}^4 : \|(i, r, I, R) - (i_c, r_c, I_c, R_c)\| \leq \delta_0\}$ be the ball of radius δ_0 center at the unperturbed solution with $0 < \delta < \delta_0$.

It is easy to check that:

Proposition 1. *The operator T is well-defined on \mathcal{B}_{δ_0} .*

The argument for the existence of the singular solution will follow directly from the Contraction Mapping Principle. By examining the definition of T we can observe that all the terms in each of its components have δ or δ^* as a coefficient. This would be fundamental in the proof that T is contraction for $0 < \delta < \delta^* \ll 1$. The following two results summarize the complete argument that we are using to establish the existence of the fixed point solution.

Proposition 2. *The T operator is a contraction on \mathcal{B}_{δ_0} for all sufficiently small $\delta > 0$.*

In order to prove Proposition 2 we need the following Lemma.

Lemma 3.1. *The growth function $\alpha(t)$ is a Lipschitz function.*

Proof. Let $\alpha(x(t), t) = (1-\delta)\rho - \frac{R_0}{c}(1-2\delta)R(t) - \frac{\delta R_0^2}{c^2\rho}R(t)^2$ be the growth function for the i function. Note that α only depends on R , so we will rewrite it as:

$$\alpha(R(t)) = (1-\delta)\rho - \frac{R_0}{c}(1-2\delta)R(t) - \frac{\delta R_0^2}{c^2\rho}R(t)^2, \quad (57)$$

Let $R_1, R_2 \in \mathcal{B}_{\delta_0}$,

$$\begin{aligned} |\alpha(R_1) - \alpha(R_2)| &\leq \frac{R_0}{c}(1-2\delta)|R_1 - R_2| + \frac{\delta R_0^2}{c^2\rho}|R_1^2 - R_2^2| \\ &= |R_1 - R_2| \left(\frac{R_0}{c}(1-2\delta) + \frac{\delta R_0^2}{c^2\rho}|R_1 + R_2| \right) \\ &\leq |R_1 - R_2|(A + B|R_1 + R_2|) \end{aligned}$$

for some constants A, B so that $A > \frac{R_0}{c}(1-2\delta)$, and $B > \frac{\delta R_0^2}{c^2\rho}$. Also note that $|R_1 + R_2|$ is bounded on \mathcal{B}_{δ_0} , thus

$$|\alpha(R_1) - \alpha(R_2)| \leq M |R_1 - R_2|. \quad (58)$$

Therefore, the growth function is Lipschitz with constant M . \square

Remark 2. As a consequence of Lemma 3.1 and to the fact that the weight function $W(u) = e^{\gamma_1 u} + e^{\gamma_2 u}$ grows exponentially at both $u = \pm\infty$, we have that

$$\left| \int_t^u (\alpha(R_1) - \alpha(R_2)) ds \right| \leq \tilde{C}\delta_0$$

Proof of Proposition 2 We will prove the required condition on T by proving that each component of T is a contraction. The first component is

$$T^{(1)}x = \frac{\delta^* R_0}{c\rho} e^{\left(\int_{t_0}^u \alpha(s) ds\right)} \int_{-\infty}^u e^{\left(-\int_{t_0}^t \alpha(s) ds\right)} i(t)r(t) dt. \quad (59)$$

Rearranging (59) so that

$$T^{(1)}x = \bar{\delta} \int_{-\infty}^u e^{\left(\int_t^u \alpha(s) ds\right)} i(t)r(t) dt,$$

where $\bar{\delta} = \delta^* \frac{R_0}{c\rho}$. Let $x_1, x_2 \in \mathbb{X}^3$

$$\begin{aligned} & \left| T^{(1)}x_1 - T^{(1)}x_2 \right| \\ & \leq \bar{\delta} \int_{-\infty}^u e^{\left(\int_t^u \alpha(x_1(s)) ds \right)} \left| i_1(t)r_1(t) - i_2(t)r_2(t) e^{\left(\int_t^u \alpha(x_2(s)) - \alpha(x_1(s)) ds \right)} \right| dt \\ & \leq \bar{\delta} \int_{-\infty}^u e^{\left(\int_t^u \alpha(x_1(s)) ds \right)} \left(|i_1(t) - i_2(t)| |r_1(t)| + |i_2(t)| |r_1(t) - r_2(t)| \right. \\ & \quad \left. + |i_2(t)| |r_2(t)| \left| 1 - e^{\left(\int_t^u \alpha(x_2(s)) - \alpha(x_1(s)) ds \right)} \right| \right) dt. \end{aligned}$$

Claim 1. $\left| 1 - e^{\left(\int_t^u \alpha(x_2(s)) - \alpha(x_1(s)) ds \right)} \right| \leq C \|R_1 - R_2\|$

Proof. Note that

$$\begin{aligned} \left| 1 - e^{\left(\int_t^u \alpha(x_2(s)) - \alpha(x_1(s)) ds \right)} \right| &= \left| e^0 - e^{\left(\int_t^u \alpha(x_2(s)) - \alpha(x_1(s)) ds \right)} \right| \\ &\leq e^{\theta u} \|R_1 - R_2\| \leq C \|R_1 - R_2\| \end{aligned}$$

for some θ between 0 and $\int_t^u \alpha(x_2(s)) - \alpha(x_1(s)) ds$. \square

Thus, going back the the proof of the proposition

$$\begin{aligned} & |i_1(t) - i_2(t)| |r_1(t)| + |i_2(t)| |r_1(t) - r_2(t)| \\ & + |i_2(t)| |r_2(t)| \left| 1 - e^{\left(\int_t^u \alpha(x_2(s)) - \alpha(x_1(s)) ds \right)} \right| \end{aligned}$$

$$\leq |i_1(t) - i_2(t)| |r_1(t)| + |i_2(t)| |r_1(t) - r_2(t)| + |i_2(t)| |r_2(t)| C \|R_1 - R_2\|.$$

So,

$$\begin{aligned} & \left| T^{(1)}x_1 - T^{(1)}x_2 \right| \\ & \leq \bar{\delta} \int_{-\infty}^u e^{f(u)} (|i_1 - i_2| |r_1| + |i_2| |r_1 - r_2| + C |i_2(t)| |r_2(t)| \|R_1 - R_2\|) dt \\ & \leq \bar{\delta} \int_{-\infty}^u e^{f(u)} \frac{1}{W(u)^2} (\|i_1 - i_2\| \|r_1\| + \|i_2\| \|r_1 - r_2\| + C \|i_2\| \|r_2\| \|R_1 - R_2\|) dt \end{aligned}$$

where $f(u) = \int_t^u \alpha(x_1(s))ds$, and $W(u) = e^{\gamma_1 u} + e^{\gamma_2 u}$ is the weight function. Observe that

$$\begin{aligned}\|r_j\| &\leq \delta_0 + M_1, \\ \|i_j\| &\leq \delta_0 + M_2.\end{aligned}$$

Thus we can define the constant,

$$\mathbb{M} = \max \{ \|r_1(t)\|, \|i_2(t)\|, C \|i_2(t)\| \|r_2(t)\| \}$$

Thus,

$$\begin{aligned}\left| T^{(1)}x_1 - T^{(1)}x_2 \right| &\leq \mathbb{M}\bar{\delta} (\|i_1(t) - i_2(t)\| + \|r_1(t) - r_2(t)\| + \|R_1 - R_2\|) \\ &\leq \delta_1 \|x_1 - x_2\|\end{aligned}$$

where $\delta_1 = \mathbb{M}\bar{\delta}$. Therefore, $T^{(1)}$ is a contraction for small δ . The second component of the operator T is

$$T^{(2)}x = \frac{1 + \rho c}{2c} \int_t^{+\infty} e^{-\frac{1}{2c} \int_t^{t_1} \beta(s)ds} i(t)dt, \quad (60)$$

where $i(s) = T^{(1)}x$. Thus, for x_1 , and $x_2 \in \mathbb{X}^4$ we have

$$\left| T^{(2)}x_1 - T^{(2)}x_2 \right| \leq M \int_t^{+\infty} \left| T^{(1)}x_1 - T^{(1)}x_2 \right| dt,$$

but we have just prove that the first component $T^{(1)}$ is a contraction. So, the last inequality becomes

$$\left| T^{(2)}x_1 - T^{(2)}x_2 \right| \leq \delta_2 \|x_1 - x_2\|.$$

for some $\delta_2 \leq \delta_1 < 1$. Now we will consider the third component of T,

$$T^{(3)}x = \int_u^{+\infty} \left(\frac{m + \delta c}{m} - \frac{\delta R_0 R(s)}{\rho m} \right) i(s)ds \quad (61)$$

where again $i(s) = T^{(1)}x$. Thus,

$$\begin{aligned}&\left| T^{(3)}x_1 - T^{(3)}x_2 \right| \\ &\leq \int_u^{+\infty} \left| \left(\frac{m + \delta c}{m} - \frac{\delta R_0 R_1(s)}{\rho m} \right) i_1(s) + \left(\frac{m + \delta c}{m} - \frac{\delta R_0 R_2(s)}{\rho m} \right) i_2(s) \right| ds \\ &= \int_u^{+\infty} \left| \left(\frac{m + \delta c}{m} - \frac{\delta R_0 R_1(s)}{\rho m} \right) T^{(1)}x_1 + \left(\frac{m + \delta c}{m} - \frac{\delta R_0 R_2(s)}{\rho m} \right) T^{(1)}x_2 \right| ds \\ &\leq \int_u^{+\infty} \left(\frac{m + \delta c}{m} \left| T^{(1)}x_1 - T^{(1)}x_2 \right| + \frac{\delta R_0}{\rho m} \left| R_1(s)T^{(1)}x_1 - R_2(s)T^{(1)}x_2 \right| \right) ds \\ &\leq \int_u^{+\infty} \frac{m + \delta c}{m} \left| T^{(1)}x_1 - T^{(1)}x_2 \right| + \frac{\delta R_0}{pm} \left| R_1(s) \right| T^{(1)}x_1 \\ &\quad - T^{(1)}x_2 \left| + \frac{\delta R_0}{pm} \left| R_1(s) - R_2(s) \right| i_2(s) \right| ds\end{aligned}$$

Notice that we have just proved above that the first two terms of the previous expression are a contraction, so as in the proof for the second component of T it suffices to check the last term. Thus,

$$\begin{aligned} & \int_u^{+\infty} \frac{m + \delta c}{m} \left| T^{(1)}x_1 - T^{(1)}x_2 \right| \\ & \quad + \frac{\delta R_0}{\rho m} |R_1(s)| \left| T^{(1)}x_1 - T^{(1)}x_2 \right| + \frac{\delta R_0}{\rho m} |R_1(s) - R_2(s)| |i_2(s)| \, ds \\ & \leq \int_u^{+\infty} \frac{m + \delta c}{m} \left| T^{(1)}x_1 - T^{(1)}x_2 \right| \\ & \quad + \frac{\delta R_0}{\rho m} \frac{\|R_1\|}{W} \left| T^{(1)}x_1 - T^{(1)}x_2 \right| + \frac{\delta R_0}{\rho m} |R_1(s) - R_2(s)| \frac{\|i_2\|}{W} \, ds. \end{aligned}$$

Thus, by considering the last term in the previous inequality we obtain that:

$$\frac{\delta R_0}{\rho m} \frac{\|i_2\|}{W} |R_1(s) - R_2(s)| \leq \frac{\delta R_0}{\rho m} \frac{\|i_2\|}{W^2} \|R_1 - R_2\|.$$

Notice that $\|i_2\| \leq M_1$ for some constant M_1 , as in the proof of Proposition 2. Also note that there is an upper bound for the weight function $\frac{1}{W^2} \leq M_2$, since the function tends to 0 at both $\pm\infty$. Thus, we have

$$\frac{\delta R_0}{\rho m} \frac{\|i_2\|}{W} |R_1(s) - R_2(s)| \leq \delta' \|R_1 - R_2\|$$

for some $\delta' \leq 1$. Thus, by choosing $\delta_3 = \min \{ \delta_1, \delta' \}$ and by the proof for the first component we obtain that

$$\left| T^{(3)}x_1 - T^{(3)}x_2 \right| \leq \delta_3 \|x_1 - x_2\|$$

To complete the proof of the proposition we just need to analyze the last component of T .

$$T^{(4)}x = \int_{-\infty}^u (\delta(1 + c\rho)i(s) + \left[\frac{2\rho m - \delta + c\rho\delta}{2\rho m} - \frac{\delta R_0}{\rho m} I(s) \right] r(s)) \, ds \quad (62)$$

It suffices to analyze the third term of (62), since the other two terms are covered in previous cases. Here we have,

$$\begin{aligned} \frac{\delta R_0}{\rho m} |I_2(t)r_2(t) - I_1(t)r_1(t)| & \leq \frac{\delta R_0}{\rho m} (|I_2(t)| |r_2(t) - r_1(t)| + |I_2(t) - I_1(t)| |r_1(t)|) \\ & \leq \delta M (\|r_2(t) - r_1(t)\| + \|I_2(t) - I_1(t)\|) \\ & \leq \delta'' \|x_1 - x_2\| \end{aligned}$$

So, we can concluded that

$$\left| T^{(4)}x_1 - T^{(4)}x_2 \right| \leq \delta_4 \|x_1 - x_2\|, \text{ for } \delta_4 \leq \min \{ \delta_1, \delta_2, \delta_3, \delta'' \} < 1.$$

Therefore, the operator T is a contraction. \square

Proof of Theorem 1.2. Since it has been shown that T is a contraction for all small $\delta > 0$ By the contraction principle we have that there is an element $x \in \mathbb{X}$ so that $Tx = x$, which shows the existence of a solution x near x_c for (52). It is clear that the same proof implies the existence of a solution x near x_c on the full slow

manifold. Then the contraction mapping argument used in the proof allows us to include higher order terms in the slow manifold (48), which prove Theorem 1.2. \square

3.4. Conclusion. Despite the simplicity of the ideas of the model considered in this paper, we can observe that very interesting mathematics arise from that model due to various factors: (1) The difference in time scales between the mutation process and the epidemiological process, (2) The nonlocal nonlinearities, and (3) The dependence of the traveling wave on the parameters. This analysis suggests that the pulse approaches 0 as $z \rightarrow \pm\infty$, but the speed of the wave depends on ϵ for $\epsilon > 0$, but small. The analysis for the solution is the result of a singular perturbation argument, and therefore no conclusion can be drawn for larger ϵ . In conclusion, we can state that the pulse solutions are robust, in the sense that they persist under small perturbation, and they look qualitatively the same as the unperturbed solution.

REFERENCES

- [1] J. W. Bebernes, C. W. Li and Y. Li, *Traveling fronts in cylinders and their stability*, Rocky Mountain J. Math., **27** (1997), 123–150.
- [2] H. Berestycki and L. Nirenberg, *Traveling fronts in cylinders*, Ann. Institute Henri Poincaré, Analysis non linéaire, **9** (1992), 497–572.
- [3] N. Fenichel, *Geometric singular perturbation theory for ordinary differential equations*, J. Diff. Equations, **31** (1979), 53–98.
- [4] P. Fife and J. B. McLeod, *The approach of solutions of nonlinear diffusions to traveling fronts solutions*, Arch. Rational Mech. Anal., **65** (1977), 355–361.
- [5] P. Fife, “Mathematical Aspects of Reacting and Diffusion Systems,” “Lecture Notes in Biomathematics,” **28**, Springer-Verlag, 1979
- [6] X-J Hou and Y. Li, *Local stability of traveling wave solutions of nonlinear reaction-diffusion equations*, Discrete and Continuous Dynamical Systems, **15** (2006), 681–701,
- [7] H. Jin, Z. Y. Li and Q. X. Ye, *Traveling wave front solutions for reaction-diffusion systems, partial differential equations, and their applications*, World Sci. Publishing, (1999), 119–129.
- [8] C. K. R. T. Jones, *Geometric singular perturbation theory*, Dynamical Systems, “Lectures Notes in Mathematics,” **1609**, Springer-Verlag, (1995), 44–118.
- [9] D. G. Kendall, *Mathematical models of the spread of infection*, Mathematics and Computer Science in Biology and Medicine, (1965), 213–225.
- [10] K. Kirchgassner, *On the nonlinear dynamics of traveling fronts*, Journal of Differential Equations, **96** (1992), 256–278.
- [11] Z. Y. Li and Q. X. Ye, *Traveling front solutions for reaction-diffusion system*, J. Partial Differential Equations, **4** (1991), 1–14.
- [12] Y. Li and Y.-P. Wu, *Stability of traveling waves with non critical speeds for double degenerate Fisher-type equations*, Discrete Contin. Dyn. Syst. Ser., *B10* (2008), 149–170.
- [13] J. Lin, V. Andraesen, R. Casagrandi and S. A. Levin, *Traveling waves in a model of influenza A drift*, Journal of Theoretical Biology, **222** (2003), 437–445.
- [14] J. Medlock and M. Kot, *Spreading disease: Integro-differential equations old and new*, Math. Biosci., **184** (2003), 201–222.
- [15] J. B. Plotkin, J. Dushoff and S. A. Levin, *Hemagglutinin sequence clusters and the antigenic evolution of influenza A virus*, Proc. Natl. Acad. Sci., **99** (2002), 6263–6268.
- [16] S. Ruan, *Spatial-temporal in nonlocal epidemiological models*, “Mathematics for Life Science and Medicine,” Springer-Verlag, (2007), 99–122.
- [17] Z.-C. Wang, W.-T. Li and S. Ruan, *Existence and stability of traveling wave fronts in reaction advection diffusion equation with nonlocal delay*, J. Diff. Equations, **238** (2007), 153–200.

- [18] A. I. Volpert, Vitaly A. Volper and Vladimir A. Volpert, *Traveling wave solutions of parabolic systems*, Translated from the Russian manuscript by James F. Heyda, "Translations of Mathematical Monographs," **140**, American Mathematical Society, Providence, RI, (1994).
- [19] S. Wiggins, *Normally hyperbolic invariant manifolds in dynamical systems*, "Applied Mathematical Sciences," **105**, Springer-Verlag, 1994.

Received November 2008; revised July 2009.

E-mail address: `jriveracruz@colgate.edu`

E-mail address: `yi-li@uiowa.edu`