

2016

# UVB Generates Microvesicle Particles via Platelet-activating Factor-receptor Signaling: A Novel Pathway by which a Skin-specific Stimulus Exerts Systemic Effects

Jeffrey B. Travers

Wright State University - Main Campus, jeffrey.travers@wright.edu

Christine M. Rapp

Wright State University - Main Campus, christine.rapp@wright.edu

Katherine E. Fahy

Wright State University - Main Campus, ahy.3@wright.edu

Christina E. Borchers

Wright State University - Main Campus, christina.borchers@wright.edu

Richard Simman

Wright State University - Main Campus, richard.simman@wright.edu

*See next page for additional authors*

Follow this and additional works at: <http://corescholar.libraries.wright.edu/ptox>

 Part of the [Chemicals and Drugs Commons](#)

## Repository Citation

Travers, J. B., Rapp, C. M., Fahy, K. E., Borchers, C. E., Simman, R., Liu, L., Chen, Y., & Bihl, J. C. (2016). UVB Generates Microvesicle Particles via Platelet-activating Factor-receptor Signaling: A Novel Pathway by which a Skin-specific Stimulus Exerts Systemic Effects. . <http://corescholar.libraries.wright.edu/ptox/95>

This Presentation is brought to you for free and open access by the Pharmacology and Toxicology at CORE Scholar. It has been accepted for inclusion in Pharmacology and Toxicology Faculty Publications by an authorized administrator of CORE Scholar. For more information, please contact [corescholar@www.libraries.wright.edu](mailto:corescholar@www.libraries.wright.edu).

---

**Authors**

Jeffrey B. Travers, Christine M. Rapp, Katherine E. Fahy, Christina E. Borchers, Richard Simman, Langni Liu, Yanfang Chen, and Ji C. Bihl

Jeffrey B. Travers<sup>1,2,3</sup> Christine M. Rapp<sup>1</sup>, Katherine E. Fahy<sup>1</sup>, Christina E. Borchers<sup>1</sup>, Richard Simman<sup>1</sup>, Langni Liu<sup>1</sup>, Yanfang Chen<sup>1</sup>, Ji C. Bihl<sup>1</sup>

Departments of Pharmacology & Toxicology<sup>1</sup> and Dermatology<sup>2</sup>, Boonshoft School of Medicine at Wright State University, Dayton VA Medical Center<sup>3</sup>, Dayton Ohio



## ABSTRACT

Though Ultraviolet B radiation (290-320 nm; UVB) exerts profound systemic effects, it is absorbed only in the epidermis. One important question in photobiology is how UVB signals are emanated from the epidermis. Microvesicle particles (MVP) are small (200-1000nm diameter) membrane-bound vesicles released by numerous cell types and can be found in the circulation. MVP can contain both nuclear and cytoplasmic components and are thought to provide a mechanism by which cells transmit signals systemically. Though various stressors have been demonstrated to generate MVP, the ability of UVB to exert this effect has not been demonstrated. Previous studies from ours and other groups have established that through its ability to act as a pro-oxidative stressor, UVB generates oxidized glycerophosphocholine (Ox-GPC) lipids with Platelet-activating Factor-receptor (PAF-R) agonist activity. Our first studies examined the ability of UVB to stimulate the release of MVP. Treatment of the human keratinocyte-derived cell line HaCaT with UVB resulted in the release of MVP. Treatment of HaCaT cells with the PAF-R agonist carbamoyl-PAF (CPAF) also generated MVP release. Preincubation of HaCaT cells with antioxidants N-acetyl cysteine and vitamin C blocked MVP release in response to UVB, but had no effect on MVP release in response to CPAF. To confirm that UVB generates MVP via PAF-R agonists, we used a human epithelial cell line with/without PAF-Rs. UVB generated MVP only in the PAF-R-expressing (KBP) not the PAF-R-negative (KBM) cells. Finally, we induced suction blisters on human skin explants and human skin in vivo and tested the ability of UVB irradiation of these blisters to induce MVP release into blister fluid. These experiments determined that UVB generated MVP in human skin. These studies describe a new pathway involving MVP generated following UVB-induced PAF agonist formation that could play an important role in transmitting systemic signals following this environmental pro-oxidative stressor.

## BACKGROUND

Platelet activating factor (PAF) is a lipid-derived 1-alkyl-2-acetyl-glycerophosphocholine, produced in response to various stimuli that produces pronounced pro-inflammatory effects. These include marked vasodilatory effects, increased vascular permeability, bronchoconstriction, and platelet activation. PAF has also been implicated in sepsis [1,2]. PAF acts by binding to a cellular heterotrimeric G-protein coupled receptor PAF-R. This receptor is found on many cell types including granulocytes, B-cells, epithelial cells (keratinocytes), and some mesenchymal cells. PAF-R activation is known to induce intracellular calcium mobilization and is expressed on keratinocytes. Recent studies have demonstrated that PAF can be produced both enzymatically and through non-enzymatic oxidation of membrane glycerophosphocholines (ox-GPCs) [3].

We and others have shown that in addition to its pro-inflammatory effects, PAF-R mediate pro-oxidative stressors including UVB-induced acute inflammation as well as delayed systemic immunosuppression [4-6]. Regarding the early acute effects of UVB, our previous studies have demonstrated that the PAF-R plays an important role in the pain associated with sunburn [6], and mediates the increased UVB sensitivity in the Xeroderma Pigmentosum Complementation group A (XPA) knock-out mouse [7].

Figure 1- Biosynthesis of PAF through enzymatic and non-enzymatic pathways





Figure 2. Picture of MVP released from a cell following stress



Microvesicle particles (MVP) are small membrane-bound vesicles released by numerous cell types and can be found in the circulation. MVP can contain both nuclear and cytoplasmic constituents and are thought to provide a mechanism by which cells transmit signals systemically [8]. MVP are 200-1000nm in diameter and differ from smaller (40-200 nm) exosomes.

Given that PAF is involved in UVB-mediated acute effects, these studies were designed to define if MVP are released from keratinocytes following UVB, and if PAF was involved in this process.

## MATERIALS & METHODS

For in vitro experiments, the human keratinocyte-derived cell line HaCaT was used. HaCaT cells were treated with UVB or the PAF-R agonist Carbamoyl-PAF (CPAF) or ethanol vehicle [3,5,7]. In some experiments HaCaT cells were preincubated for one hour with 0.5mM each of the antioxidants N-acetylcysteine and ascorbic acid before UVB/CPAF treatment. In some experiments the PAF-R-negative human epidermoid cell line KB cells transduced with functional PAF-Rs (KBP) or MSCV2.1 retroviral vector alone (KBM) were used [9]. At various times post-treatment supernatants were removed and MVP isolated using differential centrifugation and quantified using a NanoSight NS300 instrument using our previously published protocol [8,10].

For ex vivo experiments, we used skin discarded from human abdominoplasty experiments. The skin was warmed to ~37-38 degrees C and suction blisters separating epidermis from dermis were created using vacuum pumps and syringes (see Fig 5). Following blister formation, the blisters were treated with either sham or UVB or topical CPAF in ethanol:DMSO (1:9 v/v). Blister fluid was obtained and weighed in tared tubes. MVP were quantified as above.

For in vivo experiments, the volar forearm of human volunteers underwent suction blistering and blisters were treated either sham or UVB. Again, blister fluid was removed and weighed and MVP quantified as outlined above.

## RESULTS




Figure 3. Effect of UVB and CPAF on MVP release in HaCaT cells.

HaCaT cells were either A) control (0) or UVB-irradiated at various fluences or B) vehicle control or various doses of CPAF. Four hours after treatment, the supernatants were removed and MVPs quantified as outlined in Methods. C) HaCaT cells were treated with 3.6 KJ/m<sup>2</sup> UVB or 100 nM CPAF or control (CON), and were harvested 1 or 4 hrs post treatment. D). HaCaT cells were preincubated with 0.5 mM of N-acetylcysteine and 0.5 mM Vitamin C for one hour before treatment with 3.6 KJ/m<sup>2</sup> UVB or 100 nM CPAF. The supernatants were removed and MVP measured 4 hr later. The data presented are the mean ± SD. MVP numbers of duplicate values from a representative experiment of at least three performed. \*Statistically (p<0.05) significant changes from control values. NS: not statistically significant from control values.




Figure 4. Effect of UVB and CPAF on MVP release in KBP vs KBM cells. KB cells stably transduced with functional PAF-Rs (KBP) or empty vector (KBM) were treated with 100 nM CPAF or irradiated with 3.6 KJ/m<sup>2</sup> UVB or control-treated. The supernatants were harvested at 4 hrs post-treatment. The data presented are the mean ± SD. MVP numbers of duplicate values from a representative experiment from at least three performed. \*Statistically (p<0.05) significant changes from control values. NS: not statistically significant from control values.

## RESULTS Cont'd




Figure 5. Suction blister formation ex vivo & in vivo




Figure 6. UVB generates MVP formation in skin.

A). Ex vivo blisters were treated with UVB (1 or 2.5 KJ/m<sup>2</sup> or DMSO or CPAF (100 or 1000 ng). B). In vivo blisters were treated with sham or UVB (1KJ/m<sup>2</sup>). MVP tested at 2h.

## SUMMARY

- UVB generates MVP in vitro in a process involving PAF-R signaling.
- UVB also generates MVP in human skin.
- As MVP are thought to be functional signaling agents, these studies could provide a mechanism whereby UVB can generate systemic effects.

## REFERENCES

- Stafioni DM, Mohanty TM, Zimmerman GA, Prescott SM. Platelet-activating factor, a pleiotropic mediator of physiological and pathological processes. *Curr Biol* 2004; 14(14):R203-205.
- Braquet P, Touqui L, Shen Y, Vargely EB. Platelet activating Factor. *Endocrinology* 2005; 147(1):145-157.
- Matsuda GK, Johnson C, Bittencourt BS, Srinivasan SR, Moore GE, Murphy RC, Zimmerman GA, Mohanty TM, Travers JB. Ultraviolet B radiation generates platelet-activating factor-like phospholipids underlying cutaneous sunburn. *Journal of Investigative Dermatology* 2004; 122(4):644-651.
- Wallerstein JF, Ulrich SE, Nguyen DX. Platelet-activating factor, a molecular sensor for cellular damage, activates systemic nitric oxide suppression. *Journal of Neuroinflammation* 2005; 18:171-179.
- Zhang Q, Yao Y, Konger RL, Siro A, Cai S, Fukuk WE, Travers JB. UVB radiation-mediated inhibition of contact dermatitis requires immunosuppression. *Journal of Investigative Dermatology* 120(1):67-74, 2009.
- Zhang Q, Siro A, Al-Hassani M, Cai S, Fukuk WE, Travers JB, Hegarty CM. Involvement of platelet-activating factor in ultraviolet B-induced hyperalgesia. *Journal of Investigative Dermatology* 120(1):67-74, 2009.
- Yoshida K, Liu L, Wang B, Li M, Fan R, Wang S, Konger RL, Travers JB. Platelet-activating factor receptor agonists mediate xeroderma pigmentosum A photosensitivity. *Journal of Biological Chemistry* 281(12):8311-21, 2012.
- Travers JB, Eshkol H, Zhang Q, Al-Hassani M, Wang S, Konger RL, Zimmerman GA. Platelet-activating Factor: Potential Targets of Combinational Therapy for Vascular Disease. *Currrent Vascular Pharmacology* 13, 448-458, 2015.
- Travers JB, Eshkol H, Zhang Q, Al-Hassani M, Wang S, Konger RL. Regulation of UVB radiation-mediated early gene expression by the epidermal platelet-activating factor receptor. *Journal of Investigative Dermatology* 128, 455-460, 2009.
- Wang J et al. Analysis of Endothelial and Progenitor Cell Released Extracellular Microvesicles and Exosomes by Using Nanoparticle Tracking Analysis Method in Combination with Microarray and Q-dot Techniques. *PLoS One* 2016 in press.

## ACKNOWLEDGEMENTS

These studies were supported in part by grants from the NIH (R01HL062996) and Basic Science VA Merit Award