

7-9-2016

## *Bistorta amplexicaulis*: A Brief Insight to its Ethnobotany

Salma Batool

*Pir Mehr Ali Shah University of Arid Agriculture, Rawalpindi, Pakistan, salmabatoolshah@gmail.com*

Muhammad Sheeraz Ahmed

*Pir Mehr Ali Shah University of Arid Agriculture, Rawalpindi, Pakistan*

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>



Part of the [Biology Commons](#), and the [Physical Sciences and Mathematics Commons](#)

---

### Recommended Citation

Batool, S., & Ahmed, M. S. (2016). *Bistorta amplexicaulis*: A Brief Insight to its Ethnobotany, *Journal of Bioresource Management*, 3 (3).

DOI: 10.35691/JBM.5102.0056

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in *Journal of Bioresource Management* by an authorized editor of CORE Scholar. For more information, please contact [library-corescholar@wright.edu](mailto:library-corescholar@wright.edu).

## **BISTORTA AMPLEXICAULIS: A BRIEF INSIGHT TO ITS ETHNOBOTANY**

Salma Batool\* and Muhammad Sheeraz Ahmed

Department of Biochemistry, Pir Mehr Ali Shah University of Arid Agriculture Rawalpindi  
\*salmabatoolshah@gmail.com

### **Background**

*Bistorta amplexicaulis* (D. Don) Greene is an herb that belongs to the Family Polygonaceae. Its local name in Pakistan is Rain and masloon (Saqib and Sultan, 2005) and in India it is Sarbguni (Sharma *et al.*, 2004).

### **Morphology**

Plants in cultivation and in the wild are nearly always ascribable to (*Persicaria amplexicaulis*) *var. speciosa* (J.D. Hook.) characterized by its deep pinkish to reddish inflorescences (Akeroyd, 2013), and with a stem that is simple or branched, erect, with few leaves, 35-70 (to 100) cm tall, glabrous, perennial, long rhizomatous herb. The leaves are 3.0-15 x 1.75-10 cm, broadly lanceolate-ovate, serrate, acuminate, ciliate on midrib and margins, cordate or amplexicaule at base, petiole up to 10cm long. Ochrea 1.5 to 5.0 cm long, lanceolate, tubular, and acuminate with two to three long acuminate lobes. Inflorescence 1.5-10.0 cm long, terminal, simple (*var. speciosa*) or branched (*var. alba*), many flowered, dense pedunculate raceme, peduncles up to 8.0 cm long. Flowers up to 4.0 mm across, pedicel 1-3 mm long. Ochreolae 3-5 mm long, lanceolate, cartilaginous, with long aristate apex and entire margin. Tepals 5, 2-3.5 x 1.5-2.5 mm, lanceolate to ovate, obtuse, entire, unequal. Stamens 8, filaments thick and short, unequal; anthers dark bluish, subexserted to exserted. Ovary 1-2 x 0.25-0.5 (-0.75) mm, lanceolate, trigonous with three, long, filiform winged and free styles and non-prominent stigma. Nuts 3.0-5.5 x 2.0-3.5 mm, ovate, trigonous and unequal lobes, dark brown to black, glabrous, shining. Flowering period is June to

September (eflora of Pakistan). Non-glandular trichomes in the form of hair like structure, which was basally septate or not, stomata anomocytic and staurocytic (Yasmin *et al.*, 2009).

### **Distribution**

*Bistorta amplexicaulis* (syn.: *Persicaria amplexicaulis* Ronse Decraene, *Polygonum amplexicaule* (D. Don) Greene (Himalayas). A very rare and ephemeral garden escape. First seen in wasteland in Brussel in 1966 (Lambinon 1995). Recently recorded in an increasing number of localities, for instance in woodland in Beernem in 2003 and on rough ground in Schepdaal and Neder-over-Heembeek in 2011. It is abundantly found in Naran valley as well and used ethnobotanical (Khan *et al.*, 2011).

### **Ethnobotany**

In Pallas valley Pakistan, Leaves are used as a vegetable. Rhizome is used as a medicinal product. The most frequently used wild vegetables include *Allium humile*, *Kunth* (Palon), *Amaranthus hybridus* (Ganhar), *Bistorta amplexicaulis* (Rain). These plants could provide an alternative or additional income for the local people. Surveys and analyses of the degree and extent of the subsistence etc for these plants should be conducted prior to encouraging more intensive commercial exploitation (Saqib and Sultan, 2005). It is also used for making tea which is very effective in flue, fever and joints (Qureshi *et al.*, 2007)

The rhizome is applied on sores and wounds. The root is also given with the milk to women to check excess bleeding during menstruation period. Also used for

dysentery cough and tonic. Rootstock constitutes a drug Anjubar, used medicinally both in Unani and Ayurvedic system of medicine also contain tannins (Sharma *et al.*, 2004).

Its roots and leaves are highly medicinal. In Swat Kohistan, they are used for curing ulcer, rheumatic pain, backache, gout and for eyesight. Its current status is vulnerable (Hamayun *et al.*, 2006). *Bistorta amplexicaulis* is considered to purify blood (according to local traditional practitioners) and to cure ulcers in northwest Pakistan areas (Adnan and Holscher, 2010). It is also used for decoction and leaf paste are used to cure wounds, relieves dysentery and cause abortion (Bhat, 2013).

## REFERENCES

- Adnan, M. and D. Holscher. 2010. Medicinal Plant abundance in degraded and reforested sites in Northwest Pakistan. Mountain Research and Development, 30 (1): 25-32.
- Akeroyd, J. R. 2013. New nomenclatural combinations in *Persicaria*. L. Miller and a new hybrid in *Rumex* L. (*Polygonaceae*). *Contribuții Botanice*, 48: 15-21.
- Bhat, J. A., M. Kumar and R. W. Bussmann. 2013. Ecological status and traditional knowledge of medicinal plants in Kedarnath Wildlife Sanctuary of Garhwal Himalaya, India. *Journal of Ethnobiology and Ethnomedicine*, 9: 1.
- Eflora of Pakistan. [http://efloras.org/florataxon.aspx?flora\\_id=5&taxon\\_id=242100026](http://efloras.org/florataxon.aspx?flora_id=5&taxon_id=242100026)
- Hamayun, M., S. A. Khan, H. Kim, C. I. Na and I. Lee. 2006. Traditional knowledge and *ex situ* conservation of some threatened medicinal plants of Swat Kohistan, Pakistan. *International Journal of Botany*, 2 (2): 205-209.
- Khan, S. M., D. Harper, S. Page and H. Ahmad. 2011. Species and community diversity of vascular flora along Environmental gradient in Naran valley: a Multivariate Approach through indicator species analysis. *Pakistan Journal of Botany*, 43 (5): 2337-2346.
- Lambinon, J. 1995. Notes taxonomiques, nomenclaturales et chorologiques relatives à la quatrième édition de la « Nouvelle Flore » de la Belgique et des régions voisines. 3. Données nouvelles sur des plantes adventices ou subspontanées en Belgique. *Dumortiera*, 60: 1-36.
- Qureshi, R. A., M. A. Ghufuran, S. A. Gilani, K. Sultana and M. Ashraf. 2007. Ethnobotanical studies of selected Medicinal plants of sudha Gali and Ganga Chotti Hills, District Bagh, Azad Kashmir. *Pakistan Journal of Botany*, 39 (7): 2275-2283.
- Saqib, Z. and A. Sultan. 2005. Ethnobotany of Palas Valley, Pakistan. *Ethnobotanical Leaflets*, 1, Article 28.
- Sharma, P. K., N. S. Chauhan and B. Lal. 2004. Observations on traditional pytotherapy among the inhabitants of Parvati valley in Western Himalaya, India. *Journal of Ethnopharmacology*, 92: 167-176.
- Yasmin, G., M. A. Khan, N. Shaheen and M. Q. Hayat. 2009. Micromorphological Investigation of Foliar Anatomy of Genera *Aconogonon* and *Bistorta* of Family *Polygonaceae*. *International Journal of Agriculture and Biology*, 11 (3): 1814-1819.