

## Phytochemical Screening and Determination of Antibacterial, Anti-Tumorigenic and DNA Protection Ability of Root Extracts of *Saussurea Lappa*

Sughra Arif Minhas

*Institute of Natural and Management Sciences (INAM), Rawalpindi, sughra.arif@yahoo.com*

Fida Muhammad Khan

*Centre for Bioresource Research (CBR), Islamabad, bioresource123@outlook.com*

Fakhar-i- Abbas

*Centre for Bioresource Research (CBR), Islamabad, fakharabbas@hotmail.com*

Abu Ul Hassan Faiz

*Woman University of Azad Jammu & Kashmir, Bagh (AJK), sabulhussan@gmail.com*

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>



Part of the [Animal Diseases Commons](#), [Bacterial Infections and Mycoses Commons](#), [Botany Commons](#), [Immunoprophylaxis and Therapy Commons](#), [Medicinal Chemistry and Pharmaceutics Commons](#), [Other Forestry and Forest Sciences Commons](#), [Pathogenic Microbiology Commons](#), and the [Pharmacology Commons](#)

---

### Recommended Citation

Minhas, S. A., Khan, F. M., Abbas, F., & Faiz, A. H. (2017). Phytochemical Screening and Determination of Antibacterial, Anti-Tumorigenic and DNA Protection Ability of Root Extracts of *Saussurea Lappa*, *Journal of Bioresource Management*, 4 (4).

DOI: <https://doi.org/10.35691/JBM.7102.0077>

ISSN: 2309-3854 online

(Received: Sep 15, 2019; Accepted: Sep 17, 2019; Published: Sep 20, 2017)

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in *Journal of Bioresource Management* by an authorized editor of CORE Scholar. For more information, please contact [library-corescholar@wright.edu](mailto:library-corescholar@wright.edu).

---

## Phytochemical Screening and Determination of Antibacterial, Anti-Tumorigenic and DNA Protection Ability of Root Extracts of *Saussurea Lappa*

© Copyrights of all the papers published in Journal of Bioresource Management are with its publisher, Center for Bioresource Research (CBR) Islamabad, Pakistan. This permits anyone to copy, redistribute, remix, transmit and adapt the work for non-commercial purposes provided the original work and source is appropriately cited. Journal of Bioresource Management does not grant you any other rights in relation to this website or the material on this website. In other words, all other rights are reserved. For the avoidance of doubt, you must not adapt, edit, change, transform, publish, republish, distribute, redistribute, broadcast, rebroadcast or show or play in public this website or the material on this website (in any form or media) without appropriately and conspicuously citing the original work and source or Journal of Bioresource Management's prior written permission.

## PHYTOCHEMICAL SCREENING AND DETERMINATION OF ANTIBACTERIAL, ANTI-TUMORIGENIC AND DNA PROTECTION ABILITY OF ROOT EXTRACTS OF *SAUSSUREA LAPPA*

GHULAM SUGHRA ARIF<sup>1</sup>, FIDA MUHAMMAD KHAN<sup>2\*</sup>, FAKHAR-I-ABBAS<sup>2</sup> AND ABU UL HASSAN FAIZ<sup>3</sup>

<sup>1</sup>*Institute of Natural and Management Sciences (INAM), 46000 Rawalpindi, Pakistan*

<sup>2</sup>*Centre for Bioresource Research (CBR), 44000 Islamabad, Pakistan*

<sup>3</sup>*Department of Zoology, Woman University of Azad Jammu & Kashmir, 12500 Bagh, AJK, Pakistan*

\*Corresponding Email: bioresource123@outlook.com

### ABSTRACT

*Saussurea lappa*, a traditionally well-known plant has been used in different indigenous systems of medicine because of medicinal values. It is widely used in the treatment of asthma, ulcer, stomach problems and inflammatory diseases. In current study the phytochemical screening of *S. lappa* showed the presence of sesquiterpenes, anthraquinones, lignans, alkaloids, flavonoids, glycosides and steroids. Antimicrobial potential of subject plant was also investigated against three bacterial strains and maximum antibacterial activity was observed by CEE against *S. aureus*, CAE against *P. aeruginosa*, while CME, CEE, EAE and PEE showed almost same significant activity against *E. coli*. The CME (1000 ppm) was found to have maximum anti-tumor activity with 75% tumor inhibition potential. Significant DNA protection (of plasmid pBR322) activity was observed in 100 ppm CME and EAE. Therefore, *S. lappa* can be considered as promising candidate for isolation of bioactive compounds having antibacterial, antitumor and DNA protective potential.

**Keywords:** antibacterial, anti-tumor, DNA protection, *Saussurea lappa*, phytochemicals

**Abbreviations:** CME: Crude methanolic extract, CEE: Crude ethanolic extract, CAE: Crude aqueous extract, EAE: Ethyl acetate extract, PEE: Petroleum ether extract

### INTRODUCTION

Plants have been traditionally used for cancer treatment since ancient times. Isolation of potent anticancer agents including vincristine and vinblastine from *Catharanthus roseus* attracted the attention of researches for investigating anti-tumor potential of folklore plants (Cragg and Newman, 2005). Plants are also being widely used to cure infectious diseases as well. Plants are reservoir of secondary metabolites such as terpenoids, tannins, flavonoids and alkaloids which have been found to possess antimicrobial property (Cowan, 1999).

*Saussurea lappa* is the member of the family Asteraceae having 300 species (Pandey et al., 2007). Among them, *S. lappa* is a representative tall and robust

perennial herb. This specie grows on moist open slopes between the altitudes of 2600-3600 m (Zahara et al., 2014). Highest diversity is in the alpine habitats of the Himalaya and Central Asia. In Pakistan, it is mainly found in Azad Kashmir, Kaghan and Northern areas (Aswal and Mehrotra, 1994).

The medicinal value of *S. lappa* lies in some biochemical active substances that produce different physiological actions on human body. These biochemically active constituent are alkaloids, sesquiterpenes, tannins, flavonoids, phlobatanins, phenolic compounds, carbohydrates and glycosides (Chen et al., 1995; Simha et al., 2008; Sutar et al., 2011). *S. lappa* is traditionally used as aromatic, thermogenic, deodorant, digestive, carminative, anodyne,

aphrodisiac and stomachic. Stem and flowers of this plant have also been used for a long time in traditional Chinese medicines for the treatment of stomachache, dysmenorrhea, altitude sickness rheumatoid arthritis and cough with cold (Pandey et al., 2007; Sudhir et al., 2010; Madhuri et al., 2012; Lin et al., 2015).

Phytoconstituents are attributable for various pharmacological actions including anti-inflammatory, antiulcer, immunostimulant, antimicrobial, neurotoxic and cardiotoxic activity (Kulkarni, 2001; Tian and Hua, 2005; Pandey et al., 2007; Parekh and Chanda, 2007). *S. lappa* has been found to inhibit the mRNA expression of NOS (nitric oxide species) by lipopolysaccharides which stimulates the macrophages resulting in decreased production of nitric oxide (Sun et al., 2003). In the view of these aforementioned aspects current research was conducted to investigate the presence of phytoconstituents and antibacterial, anti-tumor and DNA protection potential of *Saussurea lappa*; a folklore plant.

## MATERIALS AND METHODS

### *Sample Collection*

The roots of the plant were collected from Azad Jammu and Kashmir (AJK), Pakistan and were air dried under shade at room temperature. The roots were crumbled and kept in an air tight container. Morphology of plant has been shown in Figure 1.

### *Preparation of Crude Extracts*

Crude extracts of methanol, ethanol and water were prepared. For this purpose 1.5 g of each sample was accurately weighed and dissolved in 50 mL of 100% methanol and ethanol separately. Samples were then kept in orbital shaker at 150 rpm for a week. It was followed by filtration and filtrates were kept at 40 °C allowing

the solvent to evaporate leaving behind the dried crude extracts. The crude aqueous extract was also prepared by following the aforementioned protocol.



**Figure 1:** *Saussurea lappa*; (a) Dried roots, (b) Fresh herb.

### *Preparation of Soxhlet Extract*

Two soxhlet extracts i.e., ethyl acetate and petroleum ether extracts were prepared. Extraction was done by using 200 mL of petroleum ether and ethyl acetate in soxhlet extractor. The temperature was kept less <77 °C and <90-100 °C respectively for both extracts. Samples were then collected in the collection flasks and stored at room temperature.

### *Phytochemical Analysis*

Phytochemical assays were performed for tannins, saponins, anthraquinones, amino acids, lignans, alkaloids, steroids, sesquiterpenes, glycosides and flavonoids following the published protocol (Akhtar and Farah, 1987).

### *Biological Activities*

#### *i. Antibacterial Activity*

Antibacterial activity of test samples against two gram negative (*E. coli* and *P. aeruginosa*) and one gram positive (*S. aureus*) strain was determined by following the previously used disc diffusion method with slight modifications (Nimri et al., 1999; Shafique and Qureshi,

2015). Test samples (CME, CEE, CAE, EAE and PEE) of 20 µg/mL and 40 µg/mL concentrations were used against each strain. Cefixime was employed as positive control while DMSO was employed as negative control. Zones of inhibition were measured after 24 hours.

### ii. Antitumor Assay

Anti-tumor assay was performed by following the previously illustrated protocol (Coker et al., 2003). *Agrobacterium tumefaciens* culture was used for tumor induction in potato discs. Plant extracts of 10, 100 and 1000 ppm were prepared in DMSO. Camptothecin (Sigma Chemical Co., St. Louis, MO) and DMSO were used as positive and negative controls, respectively. Culture plates were incubated at 28°C for 21 days for tumor induction. Lugol's Reagent was used for staining procedure after incubation period. Potato discs were examined using dissecting microscope and inhibition percentage was calculated the formula given below;

$$\% \text{ Inhibition} = (1 - N_s/N_c) \times 100$$

Where;

$N_s$ : Average number of tumors in sample

$N_c$ : Average number of tumors in negative control

### iii. DNA Protection Assay

DNA protection assay was performed to study the effect of plant extracts on DNA by following the previously published protocol (Tian and Hua, 2005). Each reaction was conducted in an Eppendorf tube at a total volume of 12 µl containing 3 µl of 0.5 µg plasmid DNA (pBR322, Fermentas), 3 µl of 2 mM FeSO<sub>4</sub> and 2 µl of test samples at various concentrations. After that 4 µl of 30% H<sub>2</sub>O<sub>2</sub> was added and the mixture was kept at 37 °C for 1 hr. The mixture was employed to agarose gel electrophoresis

followed by the staining of DNA bands. Quantification was done by scanning the intensity of bands with Quantity One programme (version 4.2.3, BioRad Co.).

## RESULTS

### Phytochemical evaluation

Phytochemical evaluation of various extracts of *S. lappa* revealed the presence of a wide variety of phytoconstituents as shown in Table 1.

**Table 1: Phytochemical analysis of root extracts of *S. lappa*.**

Phyto-chemicals	Results	Phyto-chemicals	Results
Alkaloids	+	Tannins	-
Lignans	+	Amino acid	+
Steroids	+	Glycosides	+
Flavonoids	+	Saponins	-
Anthraquinones	+	Sesqui-terpenes	+


(+); Presence of compounds, (-) Absence of compounds

### Biological Activities


#### i. Antibacterial Assay

Root extracts of *S. lappa* exhibited significant antibacterial activity against tested bacterial strains (*S. aureus*, *E. coli* and *P. aeruginosa*). Antibacterial potential was found to increase in direct proportion with increasing concentration of test extracts. In case of *S. aureus*, maximum activity was exhibited by CME and CEE, while CAE showed minimum antibacterial potential (Figure 2).


CAE showed highest antibacterial activity against *P. aeruginosa* while PEE was minimally active against this strain. Activities of CME, CEE and EAE were also significant (Figure 3). In case of *E. coli*, lowest bacterial inhibition potential was shown by CAE while CEE, CME, PEE and EAE showed almost same inhibition values (Figure 4).



**Figure 2: Anti-bacterial activity of various extracts of *S. lappa* against *S. aureus*.**



**Figure 3: Anti-bacterial activity of various extracts of *S. lappa* against *P. aeruginosa*.**




**Figure 4: Anti-bacterial activity of various extracts of *S. lappa* against *E. coli*.**

To determine the possible class of phytochemicals attributable for antibacterial activity we developed a correlation with different constituents that were found to be present in crude extracts of *S. lappa*. Flavonoids were found to have the best correlation ( $r=0.527$ ) with antibacterial potential of *S. lappa* extracts against *P. aeruginosa*.

## ii. Antitumor Activity


Antitumor activity of root extracts of *Saussurea lappa* was determined by potato disc assay method. For this purpose, extracts with three different concentrations were used. The maximum anti-tumor activity was shown by CME and minimum by CAE. It was observed that the inhibition of tumor increases by increasing the concentration of root extract (Figure 5). A correlation was developed between antitumor potential and phytoconstituents that were found to be present in *S. lappa* extracts. Most significant value ( $r=0.857$ ) was shown by terpenes, thus suggesting that most probable class of phytochemicals responsible for antitumor potential of *S. lappa* extracts are terpenes.



**Figure 5: Anti-tumor activity of various extracts of *S. lappa* root.**

## iii. DNA Protection Assay

It was carried out on the basis of  $H_2O_2$  and  $FeSO_4$  photolysis in order to examine the DNA protection activity of root extracts of *S. lappa*. Maximum protection activity was observed by 100ppm concentration of methanolic and ethyl acetate extracts while ethanolic, petroleum ether and aqueous extracts showed less protection potential (Figure 6).



**Figure 6: DNA protection assay. Lane M: 1kb DNA ladder, Lane 1: 100 ppm CME + 2 mM FeSO<sub>4</sub> + 30% H<sub>2</sub>O<sub>2</sub> + pBR322, Lane 2: 100 ppm CEE + 2 mM FeSO<sub>4</sub> + 30% H<sub>2</sub>O<sub>2</sub> + pBR322), Lane 3: 100 ppm EAE + 2 Mm FeSO<sub>4</sub> + 30% H<sub>2</sub>O<sub>2</sub> + pBR322), Lane 4: 100 ppm PEE + 2 Mm FeSO<sub>4</sub> + 30% H<sub>2</sub>O<sub>2</sub> + pBR322), Lane 5: 100 ppm CAE + 2 Mm FeSO<sub>4</sub> + 30% H<sub>2</sub>O<sub>2</sub> +  $\mu$ L pBR322, Lane 6: plasmid DNA + 30% H<sub>2</sub>O<sub>2</sub>, Lane 7: plasmid DNA (pBR322).**

## DISCUSSION

Since the existence of human being on earth, plants are being used for food and medicine as well. Nature has gifted plants with various potentials which are beneficial for their users. The need of the hour is to screen a number of medicinal plants for promising biological activities in order to design novel drugs for the treatment of different diseases. The phytoconstituents evaluation of the root of *S. lappa* indicated the presence of alkaloid, flavonoids, glycosides sesquiterpenes, anthraquinones, lignans and steroids. However, saponins and tannins were not detected in current study.

Phytoconstituents possess therapeutic potentials, for example, the presence of flavonoids and resins are attributable for anti-inflammatory activity (Chandur et al., 2011). Phenolics are important class of phytoconstituents including flavonoids and tannins, that possess antioxidant potentials (Okamura et al., 1993; Rice-Evans et al., 1996; Kähkönen et al., 1999). Since these compounds were found in extracts under study that might be responsible for the potent antioxidant activity of *S. lappa* in current research.

Antibacterial activity was tested against two gram-negative strains (*E. coli* and *P. aeruginosa*) and one gram positive-

strain (*S. aureus*). CEE showed maximum inhibition against the *S. aureus*, CAE showed maximum zone of inhibition against *P. aeruginosa*, while CME, CEE, EAE and PEE showed about 90% antibacterial activity against *E. coli*. It has been revealed that flavonoids can impart antibacterial impact by preventing nucleic acid synthesis and cytoplasmic membrane function (Goswami and Chatterjee, 2014; Xie et al., 2015). Flavonoids were detected in different extracts of subject plant therefore antibacterial potential of *S. lappa* might be attributed to the presence of flavonoids.

Anti-tumor activity of root extracts of *Saussurea lappa* was determined by potato disc assay method. Different concentrations of crude extracts were employed to check tumor inhibition potential that showed different percentage of tumor inhibition. Percentage inhibition of tumor was found in direct proportion to the concentration of root extracts of *S. lappa*. Twenty percent inhibition of tumor is considered as minimum significant value for anti-tumor potential of plant extract. Crude methanolic extract 1000 ppm showed maximum inhibition i.e. 75.86%. It was followed by 1000 ppm petroleum ether extract (PEE) showing 61 % inhibition, while 1000 ppm ethanolic and ethyl acetate extracts showed 54.37% and 57.82 % inhibition respectively. It is

obvious that tumor formation occurs due to imbalance between cell proliferation and cell death, which is maintained through regulation of various signal transduction pathways. Active cell death or programmed cell death (apoptosis) is caused by various physiologic or non-physiologic cell injuries including DNA damage (Folkman, 1971; Vogelstein et al., 1988). On the basis of results from preliminary studies, a cytotoxic herb *Saussurea lappa* was chosen to examine its cytotoxic effects that showed promising cytotoxic potential.

Methanolic and ethyl acetate extracts of subject plant showed promising DNA protection potential. Cells are susceptible to attack by reactive oxygen species (ROS) that are generated by both endogenous metabolisms and exogenous resources (Lenzen, 2008). Excessive production of ROS leads to oxidative stress resulting in severe pathological impacts on important bio macromolecules such as DNA and proteins. Although, cells adopt various compensatory mechanisms to counteract the deleterious impacts of oxidative stress but in some conditions, DNA damage cannot be prevented and eventually correlate with etiology, resulting in different diseases such as cancer (Birben et al., 2012). Polyphenols possess antioxidant potential and protect cells against hazardous impacts of free radicals. They can also block the action of enzymes responsible for cancer development as well as deactivate substances that promote the growth of cancers (Goswami and Chatterjee, 2014). A wide variety of polyphenols was detected in *S. lappa* that might be attributable for DNA protection potential.

## CONCLUSION

The undertaken study illustrates that *S. lappa* is an enriched reservoir of phytoconstituents possessing antitumor, antibacterial and DNA protection potential. The preliminary study further

demands bioassay guided isolation of compounds of interest from subject plant which are responsible for the aforementioned activities.

## Conflicts of Interest

The authors declare no conflict of interest

## REFERENCES

- Akhtar M, Farah N (1987). Phytochemical screening of the fruit of *Caesalpinia crista* (Karanjwa), *Melia azedarach* (Bakain) and roots of *Saussurea lappa* (Qust-e-Shireen). *Pak J Agric Sci.* 24:235-240.
- Aswal BS, Mehrotra BN, (1994). *Flora of Lahaul-Spiti (A Cold Desert in Northwest Himalaya)*. Bishen Singh Mahendra Pal Singh (Publishers and Distributors of Scientific Books), Dehra Dun, India: pp 761.
- Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O (2012). Oxidative stress and antioxidant defense. *World Allergy Organ.* 5: 9-19.
- Chandur U, Shashidhar S, Chandrasekar S, Bhanumathy M, Midhun T (2011). Phytochemical evaluation and anti-arthritic activity of root of *Saussurea lappa*. *Pharmacol.* 2: 265-267.
- Chen HC, Chou CK, Lee SD, Wang JC, Yeh SF (1995). Active compounds from *Saussurea lappa* Clarks that suppress hepatitis B virus surface antigen gene expression in human hepatoma cells. *Antivir Res.* 27: 99-109.
- Coker P, Radecke J, Guy C, Camper N (2003). Potato disc tumor induction assay: a multiple mode of drug action assay. *Phytomed.* 10: 133-138.
- Cowan MM (1999). Plant products as antimicrobial agents." *Clini Micro Reviews.* 12: 564-582.



- Cragg GM, Newman DJ (2005). Plants as a source of anti-cancer agents. *J Ethnopharma.* 100: 72-79.
- Folkman J (1971). Tumor angiogenesis: therapeutic implications. *New Engl J Med.* 285: 1182-1186.
- Goswami N, Chatterjee S (2014). Assessment of free radical scavenging potential and oxidative DNA damage preventive activity of *Trachyspermum ammi* L.(carom) and *Foeniculum vulgare* Mill.(fennel) seed extracts. *BioMed Res Intl.* 2014: 1-8.
- Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, Heinonen M (1999). Antioxidant activity of plant extracts containing phenolic compounds. *J Agric Food Chem.* 47: 3954-3962.
- Kulkarni S (2001). Immunostimulant activity of inulin isolated from *Saussurea lappa* roots. *Indian J Pharmace Sci.* 63: 292.
- Lenzen, S. (2008). Oxidative stress: the vulnerable  $\beta$ -cell. *Biochem Soc Trans.* 36: 343-347.
- Lin X, Peng Z, Su C (2015). Potential anti-cancer activities and mechanisms of costunolide and dehydrocostuslactone. *Intl J Molec Sci.* 16: 10888-10906.
- Madhuri K, Elango K, Ponnusankar S (2012). *Saussurea lappa* (Kuth root): review of its traditional uses, phytochemistry and pharmacology. *Orien Pharm Experi Med.* 12: 1-9.
- Nimri LF, Meqdam M, Alkofahi A (1999). Antibacterial activity of Jordanian medicinal plants. *Pharmace Biol* 37: 196-201.
- Okamura H, Mimura A, Yakou Y, Niwano M, Takahara Y (1993). Antioxidant activity of tannins and flavonoids in *Eucalyptus rostrata*. *Phytochem.* 33: 557-561.
- Pandey MM, Rastogi S, Rawat AKS (2007). *Saussurea costus*: botanical, chemical and pharmacological review of an ayurvedic medicinal plant. *J Ethnopharm.* 110: 379-390.
- Parekh J, Chanda S (2007). Antibacterial and phytochemical studies on twelve species of Indian medicinal plants. *Afr J Biomed Res.* 10: 175-181
- Rice-Evans CA, Miller NJ, Paganga G (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radical Bio Med.* 20: 933-956.
- Shafique N, Qureshi R (2015). Assessment of Antimicrobial and Antioxidant Activity of *Justicia californica*. *J Biore Manag.* 2: 59-62
- Simha K, Laxminarayana V, Prasad S, Khanum S (2008). Standardisation of *Yogaraja guggulu*—An Ayurvedic polyherbal formulation. *Indian J Tradit Know.* 7: 389-396
- Sudhir S, Neelima R, Barkha K, Dipika P, Baljinder K, Sarita A (2010). "Conservation of biodiversity of highly important medicinal plants of India through tissue culture technology-a review. *Agric Biol J N Am.* 1: 827-833.
- Sun CM, Syu WJ, Don MJ, Lu JJ, Lee GH (2003). Cytotoxic Sesquiterpene Lactones from the Root of *Saussurea lappa*. *J Nat Prod.* 66: 1175-1180.
- Sutar N, Garai R, Sharma US, Singh N, Roy SD (2011). Antiulcerogenic activity of *Saussurea lappa* root. *Intl J Pharm Life Sci.* 2: 516-520.
- Tian B, Hua Y (2005). Concentration-dependence of prooxidant and antioxidant effects of aloin and aloe-emodin on DNA. *Food Chem.* 91: 413-418.
- Vogelstein B, Fearon ER, Hamilton SR, Kern SE, Preisinger AC, M. Leppert M, Smits AM, Bos JL (1988). Genetic alterations during colorectal-tumor development. *New Engl J Med.* 319: 525-532.
- Xie Y, Yang W, Tang F, Chen X, Ren L (2015). Antibacterial activities of

flavonoids: structure-activity  
relationship and mechanism. *Curr*  
*Med Chem.* 22: 132-149.

Zahara K, Tabassum S, Sabir S, Arshad M,  
Qureshi RR, Amjad MS,  
Chaudhari SK (2014). A review of  
therapeutic potential of *Saussurea*  
*lappa*-An endangered plant from  
Himalaya. *Asian Pac J Trop Med.*  
7: S60-S69.