

New Record of Two Rodent Species from Khenchela Region (East of Algeria)

Hakim Drouai

University of Biskra, Biskra, Algeria, hakim.drouai@univ-biskra.dz

Yassine Noudjem

University of M'sila, M'sila, Algeria, yacinenoudjem@gmail.com

Fateh Mimeche

University of M'sila, M'sila, Algeria, fateh.mimeche@univ-msila.dz

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>



Part of the [Biodiversity Commons](#)

Recommended Citation

Drouai, H., Noudjem, Y., & Mimeche, F. (2020). New Record of Two Rodent Species from Khenchela Region (East of Algeria), *Journal of Bioresource Management*, 7 (4).

DOI: 10.35691/JBM.0202.0155

ISSN: 2309-3854 online

(Received: Nov 6, 2020; Accepted: Dec 15, 2020; Published: Dec 31, 2020)

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in *Journal of Bioresource Management* by an authorized editor of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

New Record of Two Rodent Species from Khenchela Region (East of Algeria)

Cover Page Footnote

We would like to thank Mr. Alouani Radhoune for providing the map of the studied area and Mr. Lounis Drouai for his help during the sampling.

© Copyrights of all the papers published in Journal of Bioresource Management are with its publisher, Center for Bioresource Research (CBR) Islamabad, Pakistan. Users have the right to read, download, copy, distribute, print, search, or link to the full texts of articles in the Journal. We operate under International Version 4 (CC BY 4.0) of Creative Commons Attribution License which allows the reproduction of articles free of charge with the appropriate citation of the information.

NEW RECORD OF TWO RODENT SPECIES FROM KHENCHELA REGION (EAST OF ALGERIA)

HAKIM DROUAI¹, YASSINE NOUIDJEM² AND FATEH MIMECHE^{3*}

¹Department of Agricultural Sciences, University of Biskra, Biskra, Algeria

²Department of Natural and Life Sciences, University of M'sila, M'sila, Algeria

³Department of Agricultural Sciences, University of M'sila, M'sila, Algeria

*Corresponding author: fateh.mimeche@univ-msila.dz

ABSTRACT

Mus spretus is endemic to the Mediterranean region and *Jaculus orientalis* is only found in North Africa and Saudi Arabia. Their presence in Algeria was previously evidenced in other studies, the present paper focuses on the recent discovery of the *Jaculus orientalis* Erxleben, 1777 and *Mus spretus* Lataste, 1883 in a new locality of the East of Algeria, in Khenchela region (Aurès Mountain), based on weight and external measurements. The sampling period took four months between April and July 2019. This observation will enrich the knowledge of Algerian Rodents fauna in general and of Khenchela in particular.

Keywords: Rodents fauna, Diversity, Aurès, Algeria, Mediterranean.

INTRODUCTION

Rodents are distributed over the entire world occupying all types of habitats (Hubert, 1984; Aulagnier and Thevenot, 1986; Aulagnier et al., 2017). Inventories of rodent species made in Algeria by several researchers (Kowalski and Rzebik-Kowalska, 1991; Bachar and Balhamra, 2012; Souttou et al., 2012; Beddiaf et al., 2013; Adamou et al., 2015; Hadjoudj et al., 2015; Aulagnier et al., 2017; Drouai et al., 2018) assign these to Sciuridae, Ctenodactylidae, Gliridae, Dipodidae and Muridae families. The subfamily of Dipodinae (family Dipodidae) in Algeria is represented by two species, *Jaculus jaculus* and *J.orientalis* (Ben Faleh et al., 2010; Aulagnier, 2016; Aulagnier et al., 2017).

The subfamily Murinae (family Muridae) has a distribution spanning the old world, including whole of Africa (Lecompte et al., 2008). 25% of all extant murine species live in Africa, including 32 endemic genera (Musser and Carleton, 2005). Two genera are lives between Africa and Eurasia. One of these is the

genus *Mus* (Veyrunes et al., 2005). In North Africa, *M. spretus* occupy the Maghreb area. In Algeria it appears in the Mediterranean belt and to the south it reaches the northern border of the Aures Mountains (Kowalski and Rzebik-Kowalska, 1991).

Here we present a preliminary data of new locality of *Jaculus orientalis* and *Mus spretus* in Khenchela region (semiarid region), aiming to evaluate the status of this species. We have made many sampling but this was the best region for the collection of these two species.

MATERIAL AND METHODS

Study Area

The study was carried out in Taouziant (Khenchela), located in the East of Algeria. The area has agricultural and natural landscapes in equal proportions. The average of the maximum temperature (34.9 °C) is observed in July, and the average minimum (1.85 °C) in January. The average annual precipitation is 508.83 mm, classified as semi-arid region

(Drouai, 2018). The trap sampling was carried in natural area (35°29'56.7"N and 6°45'18.8"E) (Figure 1) with an altitude of 945 m above sea line. *Artemisia herba alba* Asso., *Frangula alnus* Mill., and *Retama retam* Webb & Berthel were the main plant species at the sampling site.

Sampling

The sampling was carried out between mid-April and mid-July 2019. We used 40 wire traps of BTS type (Besançon Technique Service, France) baited with bread soaked in tuna oil (Ben Hamou et al., 2006). The traps were placed in two lines of 100 m long, 25 m apart, and 5m between each trap in each line. Traps were baited and set between 5:00 p.m. and 7:00 p.m. for four consecutive nights per season. They were checked between 7 a.m. and 9 a.m. the next day (Avenant, 1997; Hadjoudj et al., 2015; Avenant and Cavallini, 2007).




Figure 1: Map of the study area

Captures were moved alive to the laboratory, identified to species level using the identification key (Aulagnier and Thevenot, 1986; Aulagnier et al., 2008; Aulagnier et al., 2017). Then, the specimens were euthanized and standard external body measurements were recorded: body length with head (BH), length of tail (T), hind foot length (HF) and ear length (E) were also taken in mm.

RESULTS

Two species were captured in study area, which belonged to two genera and two sub-families (Table 1). The trapped species were: two specimens (adult female and young male) of *Jaculus orientalis* Erxleben, 1777, and three specimens (adult male, adult female young male) of the Algerian mouse *Mus spretus* Lataste, 1883. Their measurements are reported in Table 2.

Table 1: Rodent species collected at Taouziant (Khenchela).

Subfamily	Species	Our Study
Murinae	<i>Mus spretus</i> (Lataste, 1883)	3
Dipodinae	<i>Jaculus orientalis</i> (Erxleben, 1777)	2

DISCUSSION

According to the wild mammal catalogue of Morocco (Aulagnier and Thevenot, 1986; Aulagnier et al., 2017), these specimens correspond to *Jaculus orientalis*. These authors have reported the same morphological characters, as well as similar measurements (Table 2). Bouhaddou (2016) reported an average length of the tail of 128 to 250 mm. Kirmiz (1962) note that males are slightly larger than females and that the average body mass is 139.1g. *J. orientalis* has a distribution extending from Morocco to southern parts of Israel (Aulagnier, 2016). Biche et al., (2001) mentioned that this species is a prey of the desert grand duke *Bubo ascalaphus* in the Mergueb Nature Reserve (M'Sila, Algeria). The presence of the Greater Jerboa Egyptian in this region, which is located at the limit of the known current distribution of both: *Jaculus orientalis* (to the North of Algeria) and *Jaculus jaculus* (to the South of Algeria).

Many authors in Algeria have reported the presence of *Mus spretus* in the Tiaret region by Adamou-djerbaoui et al., (2015), in Touggourt, Southeast Algeria (Hadjoudj et al., 2015), Souttou et al., (2012) in Djelfa, Khammes et al., (2006) and Khidas et al., (1999) in Kabylie. The external measurements of this species are similar of these authors (Table 2).

Mus spretus is known to occupy a wide range of habitats in the Maghreb (crops, orchards, matorrals, juniperaies, and open forests) up to cultivated areas of oases and it reaches 1600 m altitude in Algeria and Morocco (Kowalski and Kowalska, 1991; Khidas, 1993; Aulagnier and Thévenot, 1986).

Table 2: External measurements of *Mus spretus* and *Jaculus orientalis* in this study and observed by the other authors

Species	Sex	Weight(g)	BH (mm)	T (mm)	HF (mm)	E (mm)	
<i>Jaculus orientalis</i>	Adult female	161.6	149.0	238.5	78,7	29.7	This study
	Young male	137.9	142.8	227.4	73.0	26.9	
	-	-	130-150	220-250	72-80	24-30	Aulagnier & Thevenot (1986)
<i>Mus spretus</i>	Female	19	84.4	67	17.3	13.7	
	Young male	16.5	72.3	54	15.6	11.2	This Study
	Male	20.1	86	69	18	14.4	
	-	-	70-85	50-70	15-18	11-15	Aulagnier & Thevenot (1986)
	-	18.33	80	75.33	17	9.67	Hadjoudj et al., (2015) (Mean)
-	-	76.77	57.83	16.83	12.76	Khammes et al., (2006) (Mean)	

BH: Body length with head, T: length of tail, HF: hind foot: and E: Ear lengths

CONCLUSION

Whereas the data on rodents in semiarid region stay insufficient, this work underscores that the altitude and the climate would be a combination of ecological factors in this region which largely determine the distribution and the diversity of the rodents in East Algeria.

REFERENCES

Adamou-Djerbaoui M, Labdelli F, Djelaila Y, Oulbachir K, Adamou MS and

Denys C (2015). Inventaire des rongeurs dans la région de Tiaret (Algérie). Travaux de l'Institut Scientifique, Série Générale., 8:105–112.

Aulagnier S and Thevenot M (1986). Catalogue des mammifères sauvages du Maroc. Travaux de l'Institut Scientifique, série Zoologie., 4:163.

Aulanier S, Haffner P, Mitchell-Jones T, Moutou F and Zima J (2008). Guide de mammifère d'Europe, d'Afrique

- du Nord et du Moyen-Orient. Ed. DelachauxetNiestlé, France.
- Aulagnier S (2016). *Jaculus orientalis*, Greater Egyptian Jerboa. (errata version published in 2017) The IUCN Red List of Threatened Species. Available at <http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T9123A22463965>.
- Aulagnier S, Cuzin F and Thévenot M (2017). Mammifères sauvages du Maroc. Peuplement, Répartition, Ecologie. Société française pour l'étude et la protection des mammifères, Paris: pp 339.
- Avenant NL (1997). Mammals recorded in the QwaQwa National Park (1994 - 1995). *Koedoe*, 40:31–40.
- Avenant NL and Cavallini P (2007). Correlating rodent community structure with ecological integrity, Tussen-die-Riviere Nature Reserve, Free State province, South Africa. *Integr Zool.*, 2: 212–219.
- Bachar MF and Belhamra M (2012). Contribution à l'étude de la dynamique des populations des rongeurs sauvages dans la zone de Biskra. *Courrier de Savoir*, 13:71–81.
- Beddiaf R, Sekour M, Bissati-Bouafia S, and Souttou K (2013). Inventaire des rongeurs par piégeage direct dans la région de Djanet (Tassili N'ajjer, Sahara centrale). *Revue des Bio-Ressources.*, 3(1):41–49.
- Ben Faleh A, Said K, Ben Othmen A and Granjon L (2010). Phylogeography of the Greater Egyptian Jerboa (*Jaculus orientalis*) (Rodentia: Dipodidae) in Mediterranean North Africa. *Folia Biol. (Krakow).*, 58 (3): 27–38.
- Ben Hamou M, Ben Abderrazak S, Frigui S, Chatti N and Ben Ismail R (2006). Evidence for the existence of two distinct species: *Psammomys obesus* and *Psammomys vexillaris* within the rats (Rodentia, Gerbillinae), reservoirs of cutaneous leishmaniasis in Tunisia. *Infect Genet Evol.*, 6(4): 301-308.
- Biche M, Sellami M, Libois R and Yahiaoui N (2001). Régime alimentaire de Grand-Duc de désert *Bubo ascalaphus* dans la réserve naturelle de Mergueb (M'Sila, Algérie). *Alauda.*, 69 (4): 554–557.
- Bouhaddou N (2016). Contrôle par les saisons et la photopériode de la fonction de reproduction chez la gerboise (*Jaculus orientalis*): Rôle et mécanisme d'action de la Mélatonine et du 5-Méthoxytryptophol. Doctorat Thesis, University Mohammed V, faculty of sciences, Rabat (Morocco), pp: 158.
- Drouai H, Belhamra M and Mimeche F (2018). Inventory and distribution of the rodents in Aurès Mountains and Ziban oasis (Northeast of Algeria). *An Biol.*, 40: 47–55.
- Drouai H (2018). Biodiversité des populations de Rongeurs Muridés (Rodentia: Mammalia) dans la région des Aurès et des Ziban: Approche biométrique et analytique. Doctorat Thesis, University of Biskra (Algérie), pp: 144.
- Dupuy G, Grosbety B and Dejaifve PA (2007). Suivi par piégeages de 6 espèces de micromammifères dans la Réserve Naturelle Nationale du Val d'Allier. Direction régional de l'environnement Auvergne, France.
- Hadjoudj M, Souttou K and Doumandji S (2015). Diversity and Richness of Rodent Communities in Various Landscapes of Touggourt Area (Southeast Algeria). *Acta Zool. Bulg.*, 67(3): 415–420.
- Hubert B (1984). Les rongeurs et les problèmes qu'ils posent aux cultures et aux stocks. Laboratoire de Zoologie Appliquée d'Office de la Recherche Scientifique et

- Technique Outre-Mer, Dakar (Sénégal), pp: 18
- Khammes N, Lek S and Aulagnier S (2006). Identification biométrique des deux espèces sympatriques de souris *Mus musculus domesticus* et *Mus spretus* en Kabylie du Djurdjura (Algérie). *Rev Suisse Zool.*, 113 (2): 411–419.
- Khidas K (1993). Distribution des rongeurs en Kabylie du Djurdjura (Algérie). *Mammalia.*, 57: 207–212.
- Khidas K, Khammes N and Khelloufi S (1999). Répartition spatiale et sélection de l'habitat chez le Mulot sylvestre (*Apodemus sylvaticus* L., 1758) et la Souris sauvage (*Mus pretus* Lataste, 1883) en Kabylie du Djurdjura (Algérie). *Revue Sciences et Technologie.*, 12(2): 59-64.
- Kirmiz JP (1962). Adaptation to desert environment. A study on the jerboa, rat and man, Butterworth & Co. Publishers, London.
- Kowalski K and Rzebik-Kowalska B (1991). The mammals of Algeria. Wroclaw: Polish Academy of Science, Poland, pp: 370.
- Lecompte E, Aplin K, Denys C, Catzefflis F, Chades M and Chevret P (2008). Phylogeny and biogeography of African Murinae based on mitochondrial and nuclear gene sequences, with a new tribal classification of the subfamily. *BMC Evol Biol.*, 8(1): 1-21.
- Musser GG and Carleton MD (2005). Superfamily Muroidea In: Mammal species of the world a taxonomic and geographic reference. 3rd ed, Wilson DE, Reeder DM. Baltimore, Johns Hopkins University, pp: 894–1531.
- Souttou K, Sekour M, Gouissem K, Hadjoudj M, Guezoul O, Doumandji S and Denys C (2012). Paramètres écologiques des rongeurs dans un milieu semi aride à Djelfa (Algérie). *Algérien J. Arid Environ.*, 2(2):28–41.
- Veyrunes F, Britton-Davidian J, Robinson TJ, Calvet E, Denys C and Chevret P (2005). Molecular phylogeny of the African pygmy mice, subgenus *Nannomys* (Rodentia, Murinae, Mus): Implications for chromosomal evolution. *Mol. Phylogenet Evol.*, 36(2):358–369.