

2014

A New Proof for a Result of Kingan and Lemos

Jesse Williams
Wright State University

Follow this and additional works at: https://corescholar.libraries.wright.edu/etd_all



Part of the [Physical Sciences and Mathematics Commons](#)

Repository Citation

Williams, Jesse, "A New Proof for a Result of Kingan and Lemos" (2014). *Browse all Theses and Dissertations*. 1182.

https://corescholar.libraries.wright.edu/etd_all/1182

This Thesis is brought to you for free and open access by the Theses and Dissertations at CORE Scholar. It has been accepted for inclusion in Browse all Theses and Dissertations by an authorized administrator of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

A NEW PROOF FOR A RESULT OF KINGAN AND LEMOS

A thesis submitted in partial fulfillment
of the requirements for the degree of
Master of Science

By

JESSE T. WILLIAMS

B.S., Wright State University, 2012

2014

Wright State University

WRIGHT STATE UNIVERSITY
GRADUATE SCHOOL

April 18, 2014

I HEREBY RECOMMEND THAT THE THESIS PREPARED UNDER MY SUPERVISION BY Jesse T. Williams ENTITLED A New Proof for a Result of Kingan and Lemos BE ACCEPTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF Master of Science.

Xiangqian Zhou, Ph.D.

Thesis Director

Weifu Fang, Ph.D., Chair

Department of Mathematics and Statistics

Committee on
Final Examination

Xiaoyu Liu, Ph.D.

Dan Slilaty, Ph.D.

Xiangqian Zhou, Ph.D.

Robert E. W. Fyffe, Ph.D

Vice President for Research and

Dean of the Graduate School

Abstract

Williams, Jesse T. M.S., Department of Mathematics and Statistics, Wright State University, 2014.
A New Proof for a Result of Kingan and Lemos.

The prism graph is the planar dual of $K_5 \setminus e$. Kingan and Lemos [4] proved a decomposition theorem for the class of binary matroids with no prism minor. In this paper, we present a different proof using fundamental graphs and blocking sequences.

Contents

1	Introduction	1
2	Preliminaries	2
3	Definitions of some binary matroids	5
4	A new proof for Theorem 1.2	7
	Bibliography	13

Dedicated to Dr. Neil Robertson on the occasion of his 75th birthday.

Chapter 1

Introduction

Our notation and terminology will generally follow Oxley [7] with one exception: we use $\text{si}(M)$ (resp. $\text{co}(M)$) to denote the simplification (resp. cosimplification) of a matroid M . A very interesting open problem in matroid theory is to characterize the class of binary matroids without $M(K_5)$ minor. Kingan and Lemos [4] recently obtained a strong partial result towards this where they proved a decomposition theorem for the class of binary matroids without Prism ($M^*(K_5 \setminus e)$) minor. We will give another proof of their results by using fundamental graphs and blocking sequences. Both their proof and ours make use of a result of Oxley [6] on binary matroids without P_9 or P_9^* -minors. The matroid P_9 is the 3-sum of F_7 and $M(K_4)$ with exactly one element deleted from the common triangle. The binary matroid Z_r has $2r + 1$ elements and can be represented by the binary matrix $[I_r|D]$ where D has $r + 1$ columns labeled b_1, \dots, b_r, c_r . The first r columns of D have zeros in the diagonals and ones elsewhere. The last column of D is all ones.

Theorem 1.1 (Oxley [6]) *If M is a 3-connected binary matroid with no minor isomorphic to P_9 or P_9^* , then either*

- (1) M is regular; or
- (2) M is isomorphic to $F_7, F_7^*, Z_r, Z_r^*, Z_r \setminus b_r$, or $Z_r \setminus c_r$ for some $r \geq 4$.

The next theorem is the main result in [4].

Theorem 1.2 *Let M be a 3-connected non-regular binary matroid with no $M^*(K_5 \setminus e)$ -minor. Then one of the following holds:*

- (1) M is isomorphic to $Z_r, Z_r^*, Z_r \setminus b_r$, or $Z_r \setminus c_r$ for some $r \geq 4$;
- (2) There exists a matroid $N \in \{P_9, P_9^*\}$ such that M has a 3-separation that is induced by a non-minimal 3-separation of N .
- (3) M has rank at most 5.

Chapter 2

Preliminaries

In this chapter, we present the main techniques that we will use in the proof.

Seymour's Splitter Theorem is a well-known inductive tool in proving matroid structural theorems. Recently, Kingan and Lemos obtained a stronger version which they call the "Strong Splitter Theorem".

Theorem 2.1 (Kingan and Lemos [5]) *Let N be a 3-connected proper minor of a 3-connected matroid M such that if N is a wheel or whirl, then M has no larger wheel or whirl minor, respectively. Let $m = r(M) - r(N)$. Then there exists a sequence of 3-connected matroids M_0, M_1, \dots, M_n for some integer $n \geq m$ such that*

- (1) $M_0 \cong N$ and $M_n = M$;
- (2) for $k \in \{1, 2, \dots, m\}$, $r(M_k) = r(M_{k-1}) = 1$ and $|E(M_k) - E(M_{k-1})| \leq 3$; and
- (3) for $k \in \{m+1, \dots, n\}$, $r(M_k) = r(M)$ and $|E(M_k) - E(M_{k-1})| = 1$.

Moreover, when $|E(M_k) - E(M_{k-1})| = 3$ for some $k \in \{1, 2, \dots, m\}$, $E(M_k) - E(M_{k-1})$ is a triad of M_k .

In Seymour's original version, every step is either a single-element extension or a single-element coextension, while the stronger version states that at most two consecutive single-element extensions occur in the sequence, unless the matroids involved have the same rank; when this happens, $|E(M_k) - E(M_{k-1})| = 3$ and the three new elements form a triad in M_k . We further remark that in the case $|E(M_k) - E(M_{k-1})| = 2$ for some $k \in \{1, 2, \dots, m\}$, there also exists a special triad in the matroid M_k . Assume that $M_{k-1} = M_k \setminus e/f$. Then M_k has a triad containing both e and f . (Since we may assume $M_k \setminus e$ is **not** 3-connected, as otherwise we can coextend directly from M_{k-1} ; Since $M_k \setminus e/f = M_{k-1}$ is 3-connected, f is in a series pair of $M_k \setminus e$. Now the claim follows from the fact that M_k is 3-connected.) We will use the Strong Splitter Theorem together with the remark to reduce the number of cases in our proofs.

Since we study only binary matroids, we will use fundamental graphs to represent binary matroids. Let $M = M([I_r|P])$ be a binary matroid and let B be the base of M represented by I_r . Now we label the rows of P by elements in B and label columns of P by element of $E \setminus B$. The *fundamental graph* of M with respect to B , denoted by $G_B(M)$, is the bipartite graph with bipartition $(B, E \setminus B)$ and bi-adjacency matrix P . When we draw the graph $G_B(M)$, we will use solid vertices to represent elements of B and use hollow vertices to represent elements of $E \setminus B$. Clearly by interchanging solid vertices and hollow ones, we obtain the fundamental graph $G_{E \setminus B}(M^*)$.

Minors and Separation of a binary matroid can be read off from its fundamental graphs. Note that contracting an element e in B is the same as deleting the corresponding row in the matrix P , therefore $G_{B \setminus \{e\}}(M/e)$ is obtained from $G_B(M)$ by deleting the solid vertex e ; similarly, deleting an element f in $E \setminus B$ is the same as deleting the corresponding column in the matrix P , so $G_B(M \setminus f)$ is obtained from $G_B(M)$ by deleting the hollow vertex f . To delete an element in B , or to contract

an element in $E \setminus B$, we will require the pivot operation: let (u, v) be an edge of $G_B(M)$ with $u \in B$ and $v \in E \setminus B$. Then $B' = B \triangle \{u, v\}$ is a base of M . The graph $G_{B'}(M)$ is obtained from $G_B(M)$ by the following two-step operation:

- (1) For any $b \in \text{neigh}(v) \setminus \{u\}$ and any $c \in \text{neigh}(u) \setminus \{v\}$, b and c are adjacent in $G_{B'}(M)$ if and only if they are not adjacent in $G_B(M)$; for all other pairs $b \in B$ and $e \in E \setminus B$, b and c are adjacent in $G_{B'}(M)$ if and only if b and c are adjacent in $G_B(M)$.
- (2) Exchange the labels u and v .

We call this operation pivoting on (u, v) in $G_B(M)$.

To delete an element $b \in B$ in $G_B(M)$, we pick an edge incident with b (such an edge exists provided b is not a coloop), pivot on that edge, and then delete the vertex b . Contracting an element $c \in E \setminus B$ is done similarly. Therefore, minors of M correspond to induced subgraphs of $G_B(M)$ up to pivoting operations.

A well-known fact about fundamental graphs is that the matroid is connected if and only if the fundamental graph is connected. One may also read off higher order separations of the binary matroid from its fundamental graphs through *joins*. Let G be a graph. A *1-join* in a graph G is a partition (X, Y) of $V(G)$ such that the set of edges with one end in X and the other end in Y induces a complete bipartite graph.

A *2-join* of G is a partition (X, Y) of $V(G)$ such that

- (1) there exist non-empty disjoint subsets X_1 and X_2 of X and non-empty disjoint subsets Y_1 and Y_2 of Y ;
- (2) X_i is completely joined to Y_i for $i \in \{1, 2\}$; and
- (3) there is no other edge between X and Y .

Let (X, Y) be a partition of $V(G)$ such that

- (1) there exist disjoint subsets X_1, X_2, X_3 of X and disjoint subsets Y_1, Y_2, Y_3 of Y ;
- (2) X_i is completely joined to Y_j and Y_k for $\{i, j, k\} = \{1, 2, 3\}$; and
- (3) there is no other edge between X and Y .

If none of X_i, Y_i with $i \in \{1, 2, 3\}$ is empty, then the partition (X, Y) is called a *6-join*; if exactly one of X_i, Y_i with $i \in \{1, 2, 3\}$ is empty, then the partition (X, Y) is called an *M-join*.

A *rank-2 join* is a 2-join, a 6-join, or an *M-join*.

It is not hard to see that a binary matroid M is 3-connected if and only if $G_B(M)$ is connected and has no 1-join (X, Y) with $\min(|X|, |Y|) \geq 2$. Moreover, M has an exact 3-separation if and only if $G_B(M)$ has a rank-2 join (X, Y) with $\min(|X|, |Y|) \geq 3$.

Blocking sequences were first defined in [1] and were used extensively in [2]. We will only use blocking sequences for 3-separations. For $A \subseteq E(M)$, denote $M/(A^c \cap B) \setminus (A^c \setminus B)$ by $M[A, B]$ where $A^c = E(M) \setminus A$, the complement of A . Let $N = M[X \cup Y, B]$ be a minor of M and let (X, Y) be an exact 3-separation of N . A sequence v_1, v_2, \dots, v_p of elements of $E(M) \setminus (X \cup Y)$ is a *blocking sequence* for (X, Y) with respect to B if

- (1) $(X, Y \cup \{v_1\})$ is not a 3-separation of $M[X \cup Y \cup \{v_1\}, B]$;
- (2) $(X \cup \{v_i\}, Y \cup \{v_{i+1}\})$ is not a 3-separation of $M[X \cup Y \cup \{v_i, v_{i+1}\}, B]$ for all $i \in \{1, 2, \dots, p-1\}$;
- (3) $(X \cup \{v_p\}, Y)$ is not a 3-separation of $M[X \cup Y \cup \{v_p\}, B]$; and
- (4) no proper subsequence of v_1, v_2, \dots, v_p satisfies (1), (2), and (3).

A blocking sequence alternates between elements of B and $E \setminus B$ [2]. The next theorem is also proved in [2].

Theorem 2.2 *Let $N = M[X \cup Y, B]$ be a minor of M and let (X, Y) be an exact 3-separation of N . Then (X, Y) does not induce a 3-separation in M if and only if there exists a blocking sequence for (X, Y) with respect to B .*

Let $M[A, B]$ be a minor of M . For $x \in E \setminus A$ and $e \in A$, we say x is parallel to e if x and e have the same neighbors in A or e is the only neighbor of x in A . The next lemma lists some important properties of a blocking sequence and the proofs can be found in [2, 3].

Lemma 2.3 *Let $N = M[X \cup Y, B]$ be a minor of a 3-connected binary matroid M and let (X, Y) be an exact 3-separation of N . Let v_1, v_2, \dots, v_p be a blocking sequence for (X, Y) with respect to B . Then*

- (1) *for $1 \leq i \leq j \leq p$, v_i, \dots, v_j is a blocking sequence for the 3-separation $(X \cup \{v_1, \dots, v_{i-1}\}, Y \cup \{v_{j+1}, \dots, v_p\})$ of $M[X \cup Y \cup \{v_1, \dots, v_{i-1}, v_{j+1}, \dots, v_p\}, B]$;*
- (2) *If $Y' \subset Y$ with $|Y'| \geq 3$, (X, Y') is an exact 3-separation of $M[X \cup Y', B]$, and $(X \cup \{v_p\}, Y')$ is not a 3-separation of $M[X \cup Y \cup \{v_p\}, B]$, then v_1, \dots, v_p is a blocking sequence for the 3-separation (X, Y') of $M[X \cup Y', B]$;*
- (3) *Suppose $|X| \geq 4$ and v_1 is parallel to $e \in X$ where $e \notin cl(Y)$ and $e \notin cl^*(Y)$ in N . If both e and v_1 are in B or both are in $E \setminus B$, we define $B' = B$; otherwise we define $B' = B \Delta \{e, v_1\}$. Then v_2, \dots, v_p is a blocking sequence for the 3-separation $(X \cup \{v_1\} \setminus \{e\}, Y)$ of $M[X \cup Y \cup \{v_1\} \setminus \{e\}, B']$, which is isomorphic to $M[X \cup Y, B]$.*
- (4) *For $1 \leq i \leq p$, v_i has at least one neighbor in $X \cup Y$.*

Chapter 3

Definitions of some binary matroids

In this chapter, we give the partial matrix representation and the fundamental graph of each binary matroid we will use. For those matroids whose partial matrices are clear, we will only provide the fundamental graph.




Fig. 1. $Prism / M^*(K_5 \setminus e)$




Fig. 2. P_9




Fig. 3. P_{10}




Fig. 4. \widetilde{K}_5




Fig. 5. N_{10}




Fig. 6. \widetilde{K}_{5,c_1}




Fig. 7. \widetilde{K}_{5,c_2}




Fig. 8. \widetilde{K}_{5,r_1}




Fig. 9. \widetilde{K}_{5,r_2}




Fig. 10. \widetilde{K}_{5,r_3}




Fig. 11. \widetilde{K}_{5,r_4}




Fig. 12. \widetilde{K}_{5,r_2} (Alternate graph)

Chapter 4

A new proof for Theorem 1.2

In this chapter, we present a new proof for Theorem 1.2 using fundamental graphs, blocking sequences, and the Strong Splitter Theorem.

First note that, by Theorem 1.1, it suffices to study 3-connected binary matroids with a P_9 - or a P_9^* -minor. The proof of the next two lemmas can be found in [9]. Let $X = \{1, 5, 6, 7\}$ and $Y = \{2, 3, 4, 8, 9\}$. Then (X, Y) is a non-minimal 3-separation of P_9 .

Lemma 4.1 *Let M be a 3-connected binary matroid with a P_9 -minor. If (X, Y) is not induced in M , then M must have a \widetilde{K}_5 -, an N_{10} -, or a P_{10} -minor.*

Lemma 4.2 *The matroid N_{10} is a splitter for the class of binary matroids with no minor isomorphic to \widetilde{K}_5 or \widetilde{K}_5^* .*

It follows from Lemma 4.1 and duality that if M is a 3-connected binary matroid with a P_9^* -minor and if (X, Y) is not induced in M , then M must have a \widetilde{K}_5^* -, an N_{10} -, or a P_{10} -minor.

From now on we assume that M is a 3-connected binary matroid with no $M^*(K_5 \setminus e)$ -minor. We may further assume that M has a P_9 - or a P_9^* -minor. It follows from Lemma 4.1 that either (X, Y) is induced in M , or M has a minor isomorphic to one of \widetilde{K}_5 , \widetilde{K}_5^* , N_{10} , and P_{10} . Note that the matroid P_{10} has an $M^*(K_5 \setminus e)$ -minor (to see this, delete the vertex 8 in Figure 3); also the matroid \widetilde{K}_5^* has an $M^*(K_5 \setminus e)$ -minor (simply contract 9 in the dual of Figure 4). So we have the following.

Lemma 4.3 *Let M be a 3-connected binary matroid with no $M^*(K_5 \setminus e)$ -minor. If the 3-separation (X, Y) of P_9 or of P_9^* is not induced in M , then M has a \widetilde{K}_5 - or an N_{10} -minor.*

It follows from Lemma 4.3 and 4.2 that we may assume that M has a \widetilde{K}_5 -minor. Since \widetilde{K}_5 is internally 4-connected, we will find all 3-connected extensions and coextensions of \widetilde{K}_5 . We will use $neigh(v)$ to denote the neighbor set of a vertex v . Note that, in the fundamental graph of \widetilde{K}_5 in Figure 4, 2 and 4 are symmetric to each other; so are 1 and 3. So there are at most three 3-connected single-element extensions of \widetilde{K}_5 as shown below. None of them can have an $M^*(K_5 \setminus e)$ -minor since they all have rank 4.

$neigh(v)$	Name	Remark
$\{2, 4\}$	\widetilde{K}_{5,c_1}	
$\{1, 2, 4\}$	\widetilde{K}_{5,c_2}	
$\{1, 2, 3\}$	\widetilde{K}_{5,c_3}	$\cong \widetilde{K}_{5,c_1}$

Next we will find all 3-connected coextensions of \widetilde{K}_5 . In Figure 4, we have symmetries among 5, 6, 7 and 8. Let v be the new solid vertex in a 3-connected coextension of \widetilde{K}_5 . Then v has at least two neighbors and it is not parallel to any of 1, 2, 3 or 4. So we have the following cases up to symmetry.

Case 1: $|neigh(v)| = 2$

$neigh(v)$	Prism minor?	Pivots	Deletions
$\{5, 6\}$	Yes	$(3, 10), (v, 5)$	3, 7
$\{5, 7\}$	Yes	None	9, 10
$\{5, 8\}$	No; \widetilde{K}_{5,r_1}		
$\{5, 9\}$	Yes	$(v, 5)$	7, 10
$\{5, 10\}$	Yes	$(3, 6), (v, 5)$	3, 9
$\{9, 10\}$	Yes	$(v, 10)$	6, 8

Case 2: $|neigh(v)| = 3$

$neigh(v)$	Prism minor?	Pivots	Deletions
$\{5, 6, 7\}$	Yes	$(4, 9), (v, 5)$	4, 6
$\{5, 8, 9\}$	Yes	$(4, 9), (v, 5)$	4, 6
$\{5, 7, 9\}$	Yes	None	9, 10
$\{5, 6, 10\}$	No; $\cong \widetilde{K}_{5,r_2}$		
$\{5, 8, 10\}$	Yes	$(4, 7), (v, 10)$	4, 5
$\{5, 7, 10\}$	Yes	None	9, 10
$\{5, 9, 10\}$	Yes	$(v, 10)$	5, 7

Case 3: $|neigh(v)| = 4$

$neigh(v)$	Prism minor?	Pivots	Deletions
$\{5, 6, 7, 8\}$	Yes	$(2, 5), (4, 7)$	6, 8
$\{5, 6, 7, 9\}$	Yes	$(4, 9), (v, 6)$	4, 5
$\{5, 6, 7, 10\}$	Yes	$(3, 7), (v, 5)$	3, 10
$\{5, 6, 9, 10\}$	Yes	$(3, 9), (v, 5)$	3, 6
$\{5, 7, 9, 10\}$	Yes	None	9, 10

Case 4: $|neigh(v)| = 5$ or 6

$neigh(v)$	Prism minor?	Remark
$\{6, 7, 8, 9, 10\}$	No	$\cong \widetilde{K}_{5,r_1}$
$\{5, 6, 7, 8, 9\}$	No	\widetilde{K}_{5,r_3}
$\{5, 6, 7, 8, 10\}$	No	\widetilde{K}_{5,r_2}
$\{5, 6, 7, 8, 9, 10\}$	No	\widetilde{K}_{5,r_4}

Lemma 4.4 *Let M be a 3-connected matroid with a \widetilde{K}_5 -minor and with no $M^*(K_5 \setminus e)$ -minor. If $r(M) \geq 5$, then M has a minor isomorphic to one of \widetilde{K}_{5,r_1} , \widetilde{K}_{5,r_2} , \widetilde{K}_{5,r_3} , and \widetilde{K}_{5,r_4} .*

Proof. It follows from the Strong Splitter Theorem that M has a 3-connected minor M' such that M' has a \widetilde{K}_5 -minor, $r(M') = 5$, and $11 \leq |E(M')| \leq 13$. Clearly we are done if $|E(M')| = 11$ as \widetilde{K}_{5,r_1} , \widetilde{K}_{5,r_2} , \widetilde{K}_{5,r_3} , and \widetilde{K}_{5,r_4} are all the 3-connected single-element coextensions of \widetilde{K}_5 with no $M^*(K_5 \setminus e)$ -minor. So we have the following two cases.

Case 1: $|E(M')| = 12$.

In this case M' is obtained from \widetilde{K}_5 by an extension and then a coextension. Suppose that $M' \setminus v / f \cong \widetilde{K}_5$. Then we may assume that $M' \setminus e$ is not 3-connected as otherwise M' contains one of the 3-connected coextensions of \widetilde{K}_5 as a minor and hence the lemma holds. Therefore, $M \setminus e$ is a series coextension of \widetilde{K}_5 . In term of the fundamental graph of $M \setminus e$, f either has only one neighbor, or it has the same neighbor set with another element in the base. Since M' is 3-connected and \widetilde{K}_5 has exactly two non-isomorphic extensions, we have the following subcases up to symmetry.

$neigh(v)$	$neigh(f)$	Minor	Pivots	Deletions
$\{2, 4\}$	$\{5, v\}$	Prism	$(4, 8), (f, 5)$	4, 9, 10
$\{2, 4\}$	$\{9, v\}$	Prism	(f, v)	6, 8, 10
$\{2, 4\}$	$\{10, v\}$	Prism	None	6, 8, 9
$\{2, 4\}$	$\{5, 6, 9\}$	Prism	$(f, 5)$	6, 7, 8
$\{2, 4\}$	$\{5, 8, 9, 10, v\}$	Prism	$(4, 8), (f, 10)$	4, 5, 9
$\{1, 2, 4\}$	$\{5, v\}$	Prism	$(3, 10), (f, 5)$	6, 7, 8
$\{1, 2, 4\}$	$\{6, v\}$	Prism	$(1, 5)$	8, 9, 10
$\{1, 2, 4\}$	$\{9, v\}$	Prism	$(4, 8), (f, v)$	4, 6, 10
$\{1, 2, 4\}$	$\{10, v\}$	Prism	$(4, 8)$	5, 7, 9
$\{1, 2, 4\}$	$\{5, 8, 9, 10\}$	Prism	$(1, 10), (2, 5)$	2, 8, 9
$\{1, 2, 4\}$	$\{5, 6, 9\}$	Prism	$(4, v)$	4, 6, 7
$\{1, 2, 4\}$	$\{6, 7, 9, 10, v\}$	Prism	$(4, v)$	4, 6, 7

Case 2: $|E(M')| = 13$.

In this case M' is obtained from \widetilde{K}_5 by two extensions and then a coextension and $E(M') - E(\widetilde{K}_5)$ is a triad of M' . So M' is obtained by extending twice then completing a triad using the two extended elements. Let $M'/f \setminus v, w \cong \widetilde{K}_5$. Note that in the fundamental graph of \widetilde{K}_{5,r_1} , we have symmetries among vertices 1, 2, 3, and 4. So we have the following cases up to symmetry.

$neigh(v)$	$neigh(w)$	$neigh(f)$	Minor	Pivots	Deletions
{2, 4}	{1, 2, 3}	{v, w}	Prism	(1, 5)	6, 8, 9, 10
{1, 2, 4}	{2, 3, 4}	{v, w}	Prism	(4, 8), (f, v)	5, 7, 9, 10
{1, 2, 3}	{1, 3, 4}	{v, w}	Prism	(3, 7), (f, w)	6, 8, 9, 10
{1, 2, 3}	{1, 2, 4}	{v, w}	Prism	(3, v), (4, w)	3, 4, 9, 10

□

The next lemma will help us reduce a large number of case checking in the rest of the proof.

Lemma 4.5 *Let M be a matroid obtained from \widetilde{K}_5 by two consecutive 3-connected coextensions. Then M has an $M^*(K_5 \setminus e)$ -minor.*

Proof. As we can see from the list of coextensions of \widetilde{K}_5 , there are eight ways to coextend \widetilde{K}_5 without producing an $M^*(K_5 \setminus e)$ -minor and they yield four non-isomorphic coextensions of \widetilde{K}_5 . Suppose that $M/u, v = \widetilde{K}_5$. It follows from unique representability and symmetry that it suffices to check the following cases; they all have an $M^*(K_5 \setminus e)$ -minor as shown in the table below.

$neigh(u)$	$neigh(v)$	Prism Minor?	Pivots	Deletions	Contraction
{5, 6, 7, 8, 10}	{5, 8}	Yes	(v, 5)	7, 10	2
{5, 6, 7, 8, 10}	{5, 6, 10}	Yes	(1, 10)	1, 9	3
{5, 6, 7, 8, 10}	{6, 7, 8, 9, 10}	Yes	(v, 10)	5, 7	10
{5, 6, 7, 8, 10}	{5, 6, 7, 8, 9}	Yes	(v, 8)	6, 10	8
{5, 6, 7, 8, 10}	{5, 6, 7, 8, 9, 10}	Yes	(v, 10)	6, 8	10
{5, 6, 7, 8, 9, 10}	{5, 8}	Yes	(4, 9)	4, 10	2
{5, 6, 7, 8, 9, 10}	{6, 7, 8, 9, 10}	Yes	(v, 7)	9, 10	7
{5, 6, 7, 8, 9, 10}	{5, 6, 7, 8, 9}	Yes	(u, 10), (v, 9)	6, 8	9
{5, 6, 7, 8, 9}	{5, 8}	Yes	(v, 8)	7, 9	8
{5, 6, 7, 8, 9}	{6, 7, 8, 9, 10}	Yes	(v, 7)	9, 10	7
{5, 8}	{6, 7}	Yes	(4, 9)	4, 10	2
{5, 8}	{6, 7, 8, 9, 10}	Yes	(v, 10)	6, 7	10

□

Next we list all 3-connected single-element extensions of \widetilde{K}_{5,r_1} . See Figure 8 for a fundamental graph of \widetilde{K}_{5,r_1} . Let u be the extension element. By symmetry, we have the following cases.

$neigh(u)$	Prism Minor?	Pivots	Deletions
{1, 2}	Yes	(2, u)	2, 9, 10
{1, v}	Yes	(4, 7), (v, u)	4, v, 6
{2, 4}	Yes	(4, u), (v, 5)	4, 7, 10
{2, v}	Yes	(3, 10)	5, 6, 7
{3, v}	No; $\widetilde{K}_{5,r_1,c_1}$		
{1, 2, 3}	Yes	(2, u)	2, 9, 10
{1, 2, 4}	No; $\widetilde{K}_{5,r_1,c_2}$		
{1, 3, v}	Yes	(4, 7), (v, u)	4, 5, 10
{2, 3, v}	Yes	(2, u)	2, 9, 10
{2, 4, v}	No; $\widetilde{K}_{5,r_1,c_3}$		
{2, 3, 4}	Yes	(4, u), (v, 5)	4, 7, 10
{1, 2, 3, v}	No; $\cong \widetilde{K}_{5,r_1,c_1}$		
{1, 2, 4, v}	Yes	(2, u)	2, 6, 7
{2, 3, 4, v}	No; $\widetilde{K}_{5,r_1,c_4}$		
{1, 2, 3, 4, v}	Yes	(4, u)	4, 6, 7

Now we are ready to prove the next lemma.

Lemma 4.6 *Let M be a matroid with a \widetilde{K}_{5,r_1} -minor. If M has no $M^*(K_5 \setminus e)$ -minor, then M has rank 5.*

Proof. By Lemma 4.5 and the Strong Splitter Theorem, it suffices to show that (1) every matroid obtained from \widetilde{K}_{5,r_1} by a 3-connected extension and then a 3-connected coextension will have an $M^*(K_5 \setminus e)$ -minor; (2) every matroid obtained from \widetilde{K}_{5,r_1} by two 3-connected extensions and then the coextension that forms a triad with the two extended elements will have an $M^*(K_5 \setminus e)$ -minor.

It follows from the table above that \widetilde{K}_{5,r_1} has four non-isomorphic 3-connected extensions. So there are four cases for (1). In each case we let w be the coextension element. So we have $M/w \setminus u \cong \widetilde{K}_{5,r_1}$. Again, we may assume that $M \setminus u$ is not 3-connected, that is, w is parallel to an element in $M \setminus u$. The next four tables show the single-element coextensions $\widetilde{K}_{5,r_1,c_1}$, $\widetilde{K}_{5,r_1,c_2}$, $\widetilde{K}_{5,r_1,c_3}$, and $\widetilde{K}_{5,r_1,c_4}$, respectively.

$neigh(u)$	$neigh(w)$	Prism Minor?	Pivots	Deletions	Contraction
$\{3, v\}$	$\{5, u\}$	Yes	(4, 8)	6, 7, 10	v
$\{3, v\}$	$\{6, u\}$	Yes	$(v, 5), (w, 6)$	$v, 7, 9$	4
$\{3, v\}$	$\{9, u\}$	Yes	(4, 7)	6, 8, 10	2
$\{3, v\}$	$\{10, u\}$	Yes	$(1, 10), (v, u)$	1, 5, 7	4
$\{3, v\}$	$\{5, 8, 9, 10, u\}$	Yes	$(v, u), (w, 10)$	$w, 5, 7$	u
$\{3, v\}$	$\{5, 6, 9, u\}$	Yes	(w, u)	$w, 9, 10$	u
$\{3, v\}$	$\{6, 7, 9, 10\}$	Yes	$(w, 10)$	6, 7, 8	4
$\{3, v\}$	$\{5, 8\}$	Yes	$(w, 8)$	6, 7, 10	4

$neigh(u)$	$neigh(w)$	Prism Minor?	Pivots	Deletions	Contraction
$\{1, 2, 4\}$	$\{5, u\}$	Yes	(4, 8), (w, u)	4, 9, 10	8
$\{1, 2, 4\}$	$\{6, u\}$	Yes	$(4, u), (w, 6)$	4, 7, w	u
$\{1, 2, 4\}$	$\{9, u\}$	Yes	(4, 8), (w, u)	4, 6, 10	v
$\{1, 2, 4\}$	$\{10, u\}$	Yes	$(4, 7), (w, 10)$	6, 8, 9	v
$\{1, 2, 4\}$	$\{5, 8, 9, 10\}$	Yes	$(4, u), (v, 8)$	4, 7, 9	8
$\{1, 2, 4\}$	$\{5, 6, 9\}$	Yes	$(4, u), (v, 8)$	4, 7, 9	8
$\{1, 2, 4\}$	$\{6, 7, 9, 10, u\}$	Yes	$(4, 9), (w, 10)$	6, 7, 8	v
$\{1, 2, 4\}$	$\{5, 8, u\}$	Yes	$(4, 9), (w, u)$	4, 5, 6	v

$neigh(u)$	$neigh(w)$	Prism Minor?	Pivots	Deletions	Contraction
$\{2, 4, v\}$	$\{5, u\}$	Yes	(4, 8), (w, u)	4, 9, 10	8
$\{2, 4, v\}$	$\{6, u\}$	Yes	$(v, 8)$	$v, 7, 10$	2
$\{2, 4, v\}$	$\{9, u\}$	Yes	(w, u)	6, 8, 10	v
$\{2, 4, v\}$	$\{10, u\}$	Yes	$(v, 8)$	5, 6, 7	1
$\{2, 4, v\}$	$\{5, 8, 9, 10, u\}$	Yes	(w, u)	7, 8, 10	u
$\{2, 4, v\}$	$\{5, 6, 9\}$	Yes	$(w, 6)$	7, 8, 10	6
$\{2, 4, v\}$	$\{6, 7, 9, 10, u\}$	Yes	$(w, 7)$	5, 6, 10	7
$\{2, 4, v\}$	$\{5, 8\}$	Yes	$(4, u), (w, 8)$	4, 7, 10	u

$neigh(u)$	$neigh(w)$	Prism Minor?	Pivots	Deletions	Contraction
$\{2, 3, 4, v\}$	$\{5, u\}$	Yes	$(w, 5)$	7, 9, 10	1
$\{2, 3, 4, v\}$	$\{6, u\}$	Yes	$(w, 6)$	7, 8, 9	4
$\{2, 3, 4, v\}$	$\{9, u\}$	Yes	$(4, 7), (w, u)$	4, 5, 10	v
$\{2, 3, 4, v\}$	$\{10, u\}$	Yes	$(v, 8), (w, 10)$	5, 7, 9	4
$\{2, 3, 4, v\}$	$\{5, 8, 9, 10, u\}$	Yes	$(v, u), (w, 9)$	7, 8, w	u
$\{2, 3, 4, v\}$	$\{5, 6, 9\}$	Yes	$(v, u), (w, 9)$	$v, w, 10$	u
$\{2, 3, 4, v\}$	$\{6, 7, 9, 10\}$	Yes	$(w, 10)$	5, 6, 7	4
$\{2, 3, 4, v\}$	$\{5, 8\}$	Yes	$(w, 8)$	6, 7, 9	4

Next we need to look at the matroids obtained by two extensions and then completing the triad. Suppose that $M/w \setminus u, x \cong \widetilde{K}_{5,r_1}$. By unique representability and symmetry, we have the following cases.

$neigh(u)$	$neigh(x)$	$neigh(w)$	Prism Minor?	Pivots	Deletions	Contraction
$\{3, v\}$	$\{1, 2, 3, v\}$	$\{u, x\}$	Yes	(w, u)	5, 7, 9, 10	4
$\{3, v\}$	$\{1, 2, 4\}$	$\{u, x\}$	Yes	(4, 8)	4, 6, 9, 10	v
$\{3, v\}$	$\{2, 4, v\}$	$\{u, x\}$	Yes	$(4, 9), (w, u)$	4, 6, 7, 10	v
$\{3, v\}$	$\{2, 3, 4, v\}$	$\{u, x\}$	Yes	(w, u)	5, 6, 7, 10	4
$\{1, 2, 4\}$	$\{2, 4, v\}$	$\{u, x\}$	Yes	$(4, 9), (w, u)$	4, 5, 8, 10	v
$\{1, 2, 4\}$	$\{2, 3, 4, v\}$	$\{u, x\}$	Yes	(w, u)	6, 8, 9, 10	1
$\{2, 4, v\}$	$\{2, 3, 4, v\}$	$\{u, x\}$	Yes	(w, u)	5, 6, 7, 10	4

□

Next we study matroids that have a \widetilde{K}_{5,r_2} -minor. Note that \widetilde{K}_{5,r_2} is not internally 4-connected.

Lemma 4.7 *Let M be a matroid with a \widetilde{K}_{5,r_2} -minor. If M has no $M^*(K_5 \setminus e)$ -minor, then either M is not internally 4-connected or M has rank 5.*

Proof. Suppose that M is internally 4-connected. Then there exists a blocking sequence for the non-minimal 3-separation of \widetilde{K}_{5,r_2} . Choose a blocking sequence that is as short as possible. It follows from Lemma 2.3 that such a blocking sequence must have length 1. Let a be the element in the blocking sequence. By Lemma 4.5, we may assume that $a \notin B$. Note that, there are three hidden symmetries in the fundamental graph of \widetilde{K}_{5,r_2} in Figure 12: pivot on $(1, 10)$, pivot on $(10, v)$, and pivot on $(3, 10)$. Up to these symmetries, we may assume that $\text{neigh}(a) = \{1, 2, 3, v\}$ or $\{1, 2, 3, 4, v\}$. In the former case, M has an $M^*(K_5 \setminus e)$ -minor; in the latter case, M has a \widetilde{K}_{5,r_1} -minor, so by Lemma 4.6, M has rank 5. \square

Note that by deleting vertices 6 and 8 in Figure 12, we get a matroid isomorphic to P_9^* , so the non-minimal 3-separation of \widetilde{K}_{5,r_2} is induced by the non-minimal 3-separation of P_9^* . Hence if M has \widetilde{K}_{5,r_2} -minor and a non-minimal 3-separation, then the non-minimal 3-separation of M is induced by a non-minimal 3-separation of \widetilde{K}_{5,r_2} , and so it is also induced by the non-minimal 3-separation of either P_9 or P_9^* .

Next we list all single-element extensions of \widetilde{K}_{5,r_3} . See Figure 10 for a fundamental graph of \widetilde{K}_{5,r_3} . Note that the symmetries between 1 and 3, and between 2 and 4. So there are eleven cases.

$\text{neigh}(u)$	Minor?	Pivots	Deletion(s)
$\{1, 2\}$	Prism	$(3, 7), (v, 8)$	3, 5, 9
$\{2, 4\}$	Prism	None	6, 8, 9
$\{1, v\}$	Prism	$(4, 9), (v, 5)$	$v, 6, 8$
$\{2, v\}$	Prism	$(4, 8)$	5, 6, 7
$\{1, 2, 3\}$	\widetilde{K}_{5,r_1}	$(3, 10), (v, 8)$	6
$\{1, 2, 4\}$	\widetilde{K}_{5,r_1}	$(3, 10), (4, 7), (v, 5)$	4
$\{1, 3, v\}$	\widetilde{K}_{5,r_1}	$(3, 10), (4, 7)$	6
$\{2, 4, v\}$	\widetilde{K}_{5,r_1}	$(3, 10), (4, 9), (v, 5)$	8
$\{1, 2, 3, 4\}$	Prism	$(4, 9)$	4, 6, 7
$\{1, 2, 3, v\}$	Prism	$(2, 9), (v, 8)$	5, 6, 7
$\{1, 2, 4, v\}$	Prism	$(4, 8)$	5, 7, 9

Note that all extensions of \widetilde{K}_{5,r_3} have either an $M^*(K_5 \setminus e)$ -minor, or a \widetilde{K}_{5,r_1} -minor. So the next lemma follows easily from Lemma 4.6 and the Strong Splitter Theorem.

Lemma 4.8 *Let M be a matroid with a \widetilde{K}_{5,r_3} -minor. If M has no $M^*(K_5 \setminus e)$ -minor, then M has rank 5.*

Finally we study matroids that have a minor isomorphic to \widetilde{K}_{5,r_4} . We will list all single-element extensions of \widetilde{K}_{5,r_4} . See Figure 11 for a fundamental graph of \widetilde{K}_{5,r_4} . Note that the symmetries between 1 and 3, and between 2 and 4. So there are eleven cases.

$\text{neigh}(u)$	Minor?	Pivots	Deletion
$\{1, 2\}$	\widetilde{K}_{5,r_1}	$(2, u), (3, 6)$	7
$\{1, 3\}$	\widetilde{K}_{5,r_1}	$(3, u), (4, 7)$	6
$\{1, v\}$	\widetilde{K}_{5,r_1}	$(1, u)$	1
$\{2, v\}$	\widetilde{K}_{5,r_1}	$(4, 9), (v, u)$	4
$\{2, 4\}$	\widetilde{K}_{5,r_1}	$(3, 9), (4, u), (v, 5)$	6
$\{1, 2, 3\}$	None; $\cong \widetilde{K}_{5,r_4,c_1}$		
$\{1, 2, 4\}$	None; $\cong \widetilde{K}_{5,r_4,c_1}$		
$\{2, 4, v\}$	None; $\widetilde{K}_{5,r_4,c_1}$		
$\{1, 2, 3, 4\}$	\widetilde{K}_{5,r_1}	$(1, 8), (2, 5), (3, u)$	3
$\{1, 2, 3, v\}$	\widetilde{K}_{5,r_1}	$(1, u)$	1
$\{1, 2, 4, v\}$	\widetilde{K}_{5,r_1}	$(1, u)$	1

Lemma 4.9 *Let M be a matroid with a \widetilde{K}_{5,r_4} -minor. If M has no $M^*(K_5 \setminus e)$ -minor, then M has rank 5.*

Proof. It follows from Lemmas 4.6, 4.5, and the Strong Splitter Theorem that we need to study only one extension of \widetilde{K}_{5,r_4} , namely $\widetilde{K}_{5,r_4,c_1}$ as shown in the table above.

First we look at matroids obtained from \widetilde{K}_{5,r_4} by an extension and then a coextension. Let u be the extension element and let w be the coextension element. By symmetry, we have the following cases.

$neigh(u)$	$neigh(w)$	Prism Minor?	Pivots	Deletions	Contraction
$\{2, 4, v\}$	$\{5, u\}$	Yes	$(1, 9)$	$1, 6, 8$	4
$\{2, 4, v\}$	$\{9, u\}$	Yes	(w, u)	$6, 8, 10$	v
$\{2, 4, v\}$	$\{10, u\}$	Yes	$(v, 8)$	$v, 6, 9$	8
$\{2, 4, v\}$	$\{5, 8, 9, 10, u\}$	Yes	$(w, 8)$	$5, 7, 9$	8
$\{2, 4, v\}$	$\{5, 6, 9\}$	Yes	$(w, 6)$	$5, 7, 8$	v
$\{2, 4, v\}$	$\{5, 6, 7, 8, 9, 10\}$	Yes	$(w, 10)$	$6, 8, 9$	10

Next we look at matroids obtained from \widetilde{K}_{5,r_4} by two extensions and then completing the triad. Let u and w be the two extension elements and let x the the coextension element. By symmetry, we may assume that $neigh(u) = \{2, 4, v\}$, $neigh(w) = \{1, 2, 3\}$, and $neigh(x) = \{u, w\}$. It is easy to check that this matroid has an $M^*(K_5 \setminus e)$ -minor. □

Bibliography

- [1] A. Bouchet, W.H. Cunningham, J.F. Geelen, *Principle Unimodular Skew-Symmetric Matrices*, *Combinatorica* 18 (1998) 461-486.
- [2] J.F. Geelen, A.M.H. Gerards, A. Kapoor, *the Excluded Minors for $GF(4)$ -Representable Matroids*, *J. Combin. Theory, Ser. B* 9 (2001) 247-299.
- [3] M. Halfan, *Matroid Decomposition*, Master's essay, University of Waterloo, 2002.
- [4] S.R. Kingan, M. Lemos, *a Decomposition Theorem for Binary Matroids with No Prism Minor*, *Graphs and Combinatorics*, in press.
- [5] S.R. Kingan, M. Lemos, *Strong Splitter Theorem*, *Annals of Combinatorics*, in press.
- [6] J.G. Oxley, *the Binary Matroids with No 4-wheel Minor*, *Trans. Amer. Math. Soc.* 301 (1987) 663-679.
- [7] J.G. Oxley, *Matroid Theory*, Oxford Univ. Press, New York (1992).
- [8] P.D. Seymour, *Decomposition of Regular Matroids*, *J. Combin. Theory Ser. B* 28 (1980), 305-359.
- [9] X. Zhou, *on internally 4-connected non-regular binary matroids*,