

2000

Cation-Anion-Coupled Cotransport in an Immortalized Neuronal Cell Line (C6 Glioma)

Kenneth B. E. Gagnon

Wright State University - Main Campus

Norma C. Adragna

Wright State University - Main Campus, norma.adragna@wright.edu

Robert E.W. Fyffe

Wright State University - Main Campus, robert.fyffe@wright.edu

Peter K. Lauf

Wright State University - Main Campus, peter.lauf@wright.edu

Follow this and additional works at: <http://corescholar.libraries.wright.edu/ptox>



Part of the [Chemicals and Drugs Commons](#)

Repository Citation

Gagnon, K. B., Adragna, N. C., Fyffe, R. E., & Lauf, P. K. (2000). Cation-Anion-Coupled Cotransport in an Immortalized Neuronal Cell Line (C6 Glioma). *Journal of General Physiology*, 116 (1), 20a.
<http://corescholar.libraries.wright.edu/ptox/4>

This Conference Proceeding is brought to you for free and open access by the Pharmacology and Toxicology at CORE Scholar. It has been accepted for inclusion in Pharmacology and Toxicology Faculty Publications by an authorized administrator of CORE Scholar. For more information, please contact corescholar@www.libraries.wright.edu.

of forskolin action on the Na/K pump. In a final myocyte, examined using low-[Cl⁻] (sulfamate) Tyrode's to diminish such contaminating current, forskolin reversibly enhanced strophanthidin-sensitive current at all voltages. So it appears that PKA, stimulated by forskolin, can increase Na/K pump current, but that caution must be exercised in equating cardiotoxic steroid-sensitive current with that generated by the Na/K pump. (Supported by NIH grant HL36783.)

57. **NH₄⁺ Inhibits K-Cl Cotransport in Low K Sheep Red Blood Cells** PETER K. LAUF, SUHAIL AHMED, and NORMA C. ADRAGNA, *Departments of Physiology & Biophysics, and Pharmacology & Toxicology, Wright State University, Dayton, Ohio*

K-Cl cotransport (COT) is mediated by several isoforms of KCC proteins whose cDNAs have been recently cloned. We have earlier demonstrated that, in red blood cells (RBCs), K-Cl COT carries equally K⁺ and Rb⁺, as well as Cl⁻ and Br⁻. To further characterize the selectivity of the K⁺ site, we explored whether NH₄⁺ competes with K⁺ or Rb⁺ and is transported by K-Cl COT in low K (LK) sheep (S) RBCs before and after treatment with N-ethylmaleimide (NEM), a known stimulator of K-Cl COT (Lauf and Theg, 1980. *Biochem. Biophys. Res. Commun.* 92:1422). To minimize CO₂-driven Cl/HCO₃⁻ exchange facilitating intracellular NH₄Cl formation, and hence osmotic hemolysis, LK SRBCs were pretreated with the band-3 anion transport inhibitor DIDS in either Cl⁻ or SO₄²⁻ media. Both ouabain-resistant Rb⁺ influx and K⁺ efflux were measured in either NH₄Cl or (NH₄)₂SO₄, and Cl-dependent Rb/K fluxes (K-Cl COT) were calculated from the difference of the values in the two anions. NH₄⁺ (increased by replacing Na⁺) reduced basal and NEM-stimulated Cl-dependent Rb influx by mixed-type inhibition as both V_{max} decreased and K_m increased with rising NH₄⁺ concentrations. The K_i for NH₄⁺, estimated from Rb influx measurements, was ~50 mM and found to be three times larger than the K_m for Rb⁺ in these experiments, whereas the apparent K_i for NH₄⁺-inhibited Cl-dependent K⁺ efflux was 45 mM. Experiments taking advantage of an external acid pH-induced conversion of NH₃ to NH₄⁺ were inconclusive: NEM per se modified the pH response of K-Cl COT. To test whether NH₄⁺ is actually transported by the K-Cl COT system, Cl-dependent hemolysis was measured in the presence and absence of calyculin and genistein, inhibitors of protein phosphatases and kinases, respectively, and of K-Cl COT. However, no indirect evidence for NH₄⁺ transport by the system could be detected with this method. The flux data suggest that in DIDS-treated LK SRBCs, NH₄⁺ binds to an external low-affinity site in the transporter and causes allosteric inhibition of K-Cl COT. These results are consistent with the lack of NH₄⁺ transport by the hemolysis approach.

58. **Cation-Anion-coupled Cotransport in an Immortalized Neuronal Cell Line (C6 Glioma)** KENNETH B.E. GAGNON,* NORMA C. ADRAGNA,† ROBERT E.W. FYFFE,§ and PETER K. LAUF,* **Department of Physiology and Biophysics, †Department of Pharmacology and Toxicology, §Department of Anatomy, and ¶Department of Brain Research Center, Wright State University, Dayton, Ohio*

Ion gradients across the cell membrane are vitally important for proper cell-to-cell communication and general cell homeostasis and are maintained by both primary (ATP-dependent) and

secondary active transport mechanisms. Among the latter, Na-K-2Cl cotransport (COT) is present in primary cultures of rat astrocytes (Tas et al. 1987. *Biochim. Biophys. Acta.* 903:411-416) and implicated in the K spatial buffering capacity of glial cells. However, no reports on the presence of glial K-Cl COT are available yet. The present work was designed to study Na-K-2Cl COT and K-Cl COT in C6 glioma cells. Cell cultures were grown to confluence under 5% CO₂ supplemented air atmosphere in 12-well plates using F12K essential medium containing 15% horse serum, 2.5% fetal bovine serum, and penicillin/streptomycin. Influx of Rb (a K congener) was measured under initial velocity conditions at 6 min and 37°C in Na media containing (mM): 10 RbCl, 2 CaCl₂, 1 MgCl₂, 10 glucose, buffered to pH 7.4 with 20 HEPES/TRIS. Rb influx (nmol/mg protein × min) was 16.0 in the absence of inhibitors, 14.3 with 1.0 mM ouabain, 7.9 with ouabain and 0.005 mM bumetanide, and 4.3 with the additional presence of 2 mM furosemide. Thus, the Na/K pump represents 10% of the total Rb influx, whereas the Na-K-2Cl COT and K-Cl COT represent 40 and 22%, respectively. Na-K-2Cl COT was inhibited in a dose-dependent manner by bumetanide, whereas 100% inhibition of K-Cl COT occurred at 2 mM furosemide. At its maximum effective concentration, NEM stimulated the Na/K pump and K-Cl COT by 2- and 10-fold, respectively, and inhibited Na-K-2Cl COT by 100%. Cell K contents remained relatively unchanged, but fell by ~20% after NEM treatment, suggesting opening of a K channel, reported by us in other cells. Using a polyclonal anti-rabbit KCC1 antibody (generously supplied by Dr. Eric Delpire), K-Cl COT in C6 glioma cells was immunohistochemically demonstrated by a strong, punctuate immunofluorescent labeling throughout the cytoplasm and cell membrane. Our results on cultured C6 glioma cells corroborate the immunological findings of Plotkin et al. (1997. *Am. J. Physiol. Cell Physiol.* 272:C173-C183) and the RT-PCR results of Payne et al. (1996. *J. Biol. Chem.* 271:16245-16252) showing the KCC1 isoform in astrocytes. Western blots using the same anti-rabbit KCC1 antibody revealed a single strong band of ~120 kD, presumably the cytosolic, unglycosylated K-Cl cotransporter. These studies have therefore established: (a) the functional presence of the Na/K pump, and both Na-K-2Cl and K-Cl COT; (b) the immunohistochemical presence in C6 glioma cells of a KCC isoform of K-Cl COT; and (c) inactivation of Na-K-2Cl COT and simultaneous stimulation of K-Cl COT by NEM, possibly implicating a shared regulatory pathway. Studies in progress are aimed at the Cl dependence and the critical NEM concentration inversely affecting Na-K-2Cl and K-Cl COT. (Supported in part by NIH, AHA, and a WSU SOM alpha grant.)

59. **Transient Nature of the Stimulatory "NEM-Effect" on K-Cl Cotransport in KCC1-transfected HEK293 and Primary Rat Aortic Smooth Muscle Cells** PETER K. LAUF,* JIN ZHANG, JING ZHANG, and NORMA C. ADRAGNA,† **Department of Physiology and Biophysics, and †Department of Pharmacology and Toxicology, Wright State University, Dayton, Ohio*

K-Cl cotransport (COT) is mediated by several isoforms of KCC proteins whose cDNAs have been recently cloned. The thiol reagent N-ethylmaleimide (NEM) (Lauf and Theg, 1980. *Biochem. Biophys. Res. Commun.* 92:1492) has been widely used to demonstrate, by several-fold stimulation, the presence of K-Cl COT in a variety of cells such as red blood cells (RBCs), rat aortic smooth muscle cells (RASMCs), and human embryonic kidney (HEK293) cells transfected with KCC isoforms. In RBCs, NEM